

Data Paper

Unpublished Mediterranean and Black Sea records of marine alien, cryptogenic, and neonative species (III)

Jennifer C.A. Pistevos¹, Argyro Zenetos², Yolanda Stranga¹, Adrian Teaca³, Tatiana Begun³, Domen Trkov⁴, Joachim Langeneck⁵, Mihaela Muresan³, Robert Comas-González⁶, Juan Sempere-Valverde^{7,8}, Andrea Toso^{5,9}, Sahar Chebaane^{7,10}, Carlos Navarro-Barranco⁸, Antonietta Rosso¹¹, Lydia Png-Gonzalez⁶, Sylvaine Giakoumi¹², Maite Vázquez-Luis⁶, José Manuel Cañizares-González⁶, Chiara Siddiolo^{11,13}, Gemma Donato¹¹, Salvatore Giacobbe¹⁴, Aina Carbonell⁶, Rym Zakhama-Sraieb^{15,16}, Emanuela Di Martino¹¹, Izdihar Ali Ammar¹⁷, Andrea Cosentino¹⁸, Intissar Mnasri Afifi¹⁶, Rossana Sanfilippo¹¹, Ghazi Bitar¹⁹, Eleni Kytinou^{1,20}, Gokhan Tari²¹, Mehmet Fatih Huseyinoglu²¹, Vasilis Gerakaris²⁰, Yiannis Issaris²⁰, Polytimi-Ioli Lardi^{1,20}, George Oikonomidis²⁰, Maria Salomidis²⁰, Konstantinos Tsirintanis²⁰, Periklis Kleitou²², George Constantinou²², Christina Michail²², Michel Bariche²³, Alan Deidun²⁴, Alessio Marrone²⁴, Michail Ragkousis¹, Manuela Falautano²⁵, Luca Castriota²⁵, Carla Morri²⁶, Carlo Nike Bianchi²⁶, Grigorios Skouradakis²⁷, Fiona Tomas Nash²⁸, Markos Digenis²⁹, Emanuele Mancini^{11,30}, Elitsa Slaveva Stefanova¹³¹, Adriana Vella³², Noel Vella³², Vesselina Vasileva Mihneva³³, Evangelos Papadimitriou¹, Marius Skolka³⁴, Kremena Stefanova³¹, Ana Fortić⁴, Maria Garcia³⁵, Thanos Dailianis²⁷, Laura Tamburello¹², Victor Surugiu³⁶, Bessy Stancanelli³⁷, Vincenzo Di Martino³⁸, Angeliki Bosmali¹, Borut Mavrič⁴, Vasilis Gerovasileiou^{27,29}, Demetris Kletou^{22,39}, Fabio Crocetta^{40,41}, Ioannis Rallis^{27,29}, Julian Evans⁴², Maria Altamirano⁴³, Leda Liyue Cai⁴⁴, Ergün Taşkın⁴⁵, Giulio Franzitta¹², Yassine Ramzi Sghaier⁴⁶, Abigail Kennedy⁴⁷, Angela G. Bartolo⁴⁸, Frithjof C. Küpper^{47,49}, Maria Rousou⁵⁰, Víctor Orenes-Salazar⁵¹, Andrea Bonifazi⁵², Louis Hadjioannou⁴⁴, Zois Vlantaridis²⁹, Hany El Shaer⁵³, Mülkibar Çiftçioğlu⁵⁴, Hasan Cerim⁵⁴, Lovrenc Lipej⁴, Daniele Grech⁵⁵, Maria Zotou¹, Ernesto Azzurro⁵⁶, Patrick J. Schembri⁴², Francesco Tiralongo^{11,57}, Georgios Chatzigeorgiou²⁷, Manfredi Di Lorenzo¹², Maria Vigo Fernandez¹², Antonio Di Franco¹², Martina Orlando-Bonaca⁴, Antonio Calò⁵⁸, Christina Malefaki⁵⁹, Jose Antonio García-Charton⁵¹, Orestis Papadakis¹, Sercan Yapıcı⁶⁰, Valentina Tanduo⁴⁰, Ilaria D'Aniello⁶¹, Marco Munari^{40,62}, Thodoros E. Kampouris^{63,64}, Matteo Putignano^{5,9}, Riccardo Virgili^{40,41,65}, Sandra Agius Darmanin⁶⁶, Furkan Bilgiç⁶⁷, Giacomo Sardo⁶⁸, Monica Calabrò⁶⁸, Athanasios Nikolaou¹, Zinovia Erga⁶⁴, Ali Zogheib⁶⁹, Paolo Balistreri⁷⁰, Senem Çağlar⁷¹, Ferhat Yalçın⁷², David Osca⁷³, Eirini Gratsia^{27,59}, Giorgio Aglieri¹², Melih Ertan Çinar⁷⁴, Mirko Mutalipassi⁴⁰, Salvatore Coco⁷⁵, Andromachi Gkoulia⁴⁴, Alfonso A. Ramos-Esplà⁷⁶, Anna Maria Mannino⁷⁷, Desirée Dimichele^{5,9,78}, Simone Spinelli⁴, Elisa Arroyo Martínez⁷⁹, Andrea Tursi⁸⁰, Annalisa Capasso⁸¹, Georgios Christidis⁸², Michele Solca^{9,134}, Mouna Rifi⁸³, Radhouan Belgacem El Zrelli⁸⁴, Ramla Bouhlel¹⁰, Riccardo Scalera⁸⁵, Ada Bandi⁸⁶, Anna Reboa⁸⁶, Antonios Mazaris⁸⁷, Hocein Bazairi^{88,89}, Marco Bertolino⁸⁶, Mohamed Selfati⁹⁰, Benedetto Sirchia⁹¹, Constantinos G. Georgiadis⁹², Donat Petricioli⁹³, Eleonora Macaluso⁹¹, Enrico D'Aniello⁹⁴, Nikolaos Fotiadis⁸², Rifat Tezel⁵⁴, Slavica Petović⁹⁵, Tatjana Bakran-Petricioli⁹⁶, Valentina Pitacco⁴, Alberto Colletti^{41,97}, Bilel Bensari⁹⁸, Lamia Bahbah^{98,99}, Salvatore D'Aniello¹⁰⁰, Spiros Psailas¹⁰¹, Stefanos Kalogirou¹⁰², Tarik Kaikai⁸⁸, Aitor Forcada¹⁰³, Alessandro Manfredonia⁴⁰, Antonella Bottalico⁸⁰, Carlos Valle¹⁰³, Coline Bedoret¹⁰⁴, Dimitrios Damalas¹⁰⁵, Domenico Sacco^{106,107}, Ernesto Mollo¹⁰⁸, Foteini Iatrou¹⁰², Frederique Cerqueira¹⁰⁹, Frédérique Viard¹⁰⁹, Guido Villani¹¹⁰, Mathieu Foulquié^{111,112}, Michele Luca Geraci⁶⁸, Miljan Šiško⁴, Oscar Ocaña¹¹³, Renaud Dupuy de la Grandrive¹¹⁴, Sergio Vitale⁶⁸, Tullio Rossi¹¹⁵, Ziad Samaha¹¹⁶, Ricardo Aguilar¹¹⁷, Helena Alvarez¹¹⁷, Selma Menabit³, Ana Bianca Pavel³, Annalisa Azzola⁸⁶, Arthur R. Bos¹¹⁸, Athanasios Anastasiadis¹¹⁹, Jorge Blanco¹¹⁷, Christophe Lejeune¹⁰⁴, Daniela Coppola¹²⁰, Dominique Davoult¹²¹, Daniel Faget¹²², Danilo Scannella¹²³, Erica Keppel¹²⁴, Fabio Falsone⁶⁸, Fortunato Palma Esposito¹²⁰, Silvia García¹¹⁷, Giulia Furfaro^{5,9,41}, Helmut Zibrowius¹²⁵, Ibrahim Iziani¹²⁶, Iro Anastasopoulou¹⁰², Konstantinos Kavakakis¹⁰², Leonardo Scavino¹²⁷, Lorenzo Merotto¹²⁸, Maria Yankova³¹, Monica Montefalcone^{41,86}, Neža Leban⁴, Nicola Ungaro¹²⁹, Paolo Mariottini¹³⁰, Allison L. Perry¹¹⁷, Petros Stavrou¹³¹, Valerio Mazzella^{41,132}, Violin Raykov¹³³, Jasmine Ferrario⁷⁵, Caterina Stamouli⁸², Apostolos Siapatis⁸², Emmanouela Vernadou²⁷, Claudia Bommarito¹², Giacomo Milisenda¹², Emanuele Somma¹², Stefania Russo¹², Stelios Katsanevakis¹

¹Department of Marine Sciences, University of the Aegean, Lofos Panepistimiou, 81100 Mytilene, Greece; ²Institute of Marine Biological Resources and Inland Waters, Hellenic Centre for Marine Research (HCMR), 19013 Anavissos, Greece; ³National Institute for Research and Development on Marine Geology and Geoecology - GeoEcoMar, Bucharest, Romania; ⁴Marine Biology Station Piran, National Institute of Biology, Fornace 41, 6330 Piran, Slovenia; ⁵Consorzio Nazionale Interuniversitario per le Scienze del Mare (CoNISMa), U.L.R. di Lecce, Campus Ecotekne, strada provinciale Lecce-Monteroni, 73100, Lecce, Italy; ⁶Instituto Español de Oceanografía (IEO-CSIC). Centro Oceanográfico de Baleares, Muelle de poniente s/n, 07015, Palma de Mallorca, Spain; ⁷Biological and Environmental Sciences and Engineering (BESE), Red Sea Research Center (RSRC), King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia; ⁸Laboratorio de Biología Marina, Facultad de Biología, Universidad de Sevilla, Avenida Reina Mercedes, Sevilla, 41012, Spain; ⁹Department of Biological and Environmental Sciences and Technologies (DiSTeBA) - Via Monteroni, 165 - 73100 Lecce, Italy - University of Salento; ¹⁰Association Notre Grand Bleu (NGB, NGO), QRHM+566, Résidence Kairouan Cap marina Monastir, Monastir 5000, Tunisia; ¹¹Department of Biological, Geological and Environmental Sciences, University of Catania, Corso Italia, 57 - 95129 and Via Androne, 81 - 95124, Catania, Italy; ¹²Stazione Zoologica Anton Dohrn, Department of Integrative Marine Ecology (EMI), Sicily Marine Center, Lungomare Cristoforo Colombo, 4521, 90149, Palermo, Italy; ¹³University School for Advanced Studies IUSS Pavia, Pavia, Italy; ¹⁴Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, Via Stagno d'Alcontres 31 - 98166 Messina, Italy; ¹⁵University of Manouba, High Institute of Biotechnology of Sidi Thabet, BiotechPôle, BP-66, 2020 Sidi Thabet, Ariana, Tunisia; ¹⁶University of Tunis El Manar, Faculty of Sciences of Tunis, Laboratory of Diversity, Management and Conservation of Biological Systems, LR18ES06, Tunis, Tunisia; ¹⁷Department of Marine Biology, High Institute of Marine Research, Latakia University, Latakia, Syria; ¹⁸ANISN, Associazione Nazionale Insegnanti Scienze Naturali, Naples, Italy; ¹⁹Department of Marine Biology, Faculty of Sciences, Lebanese University, Hadath, Beirut, Lebanon; ²⁰Hellenic Centre for Marine Research (HCMR), Institute of Oceanography (IO), Anavissos 19013, Attica, Greece; ²¹University of Kyrenia, Cyprus; ²²Marine and Environmental Research (MER) Lab, 4533, Limassol, Cyprus; ²³Department of Biology, FAS, American University of Beirut, Lebanon; ²⁴Oceanography Malta Research Group, Department of Geosciences, University of Malta, Msida, Malta MSD 2080, Malta; ²⁵Italian Institute for Environmental Protection and Research (ISPRA), Lungomare Cristoforo Colombo n.4521, Località Addaura, 90149 Palermo, Italy; ²⁶Department of Integrative Marine Ecology (EMI), Stazione Zoologica Anton Dohrn - National Institute of Marine Biology, Ecology and Biotechnology, Genoa Marine Centre (GMC), Villa del Principe, Piazza del Principe 4, 16126 Genova, Italy; ²⁷Hellenic Centre for Marine Research (HCMR), Institute of Marine Biology, Biotechnology and Aquaculture (IMBBC), 71500 Heraklion, Greece; ²⁸Instituto Mediterráneo de Estudios Avanzados (UIB-CSIC), C/ Miquel Marqués 21, 07190, Esporles, Islas Baleares, Spain; ²⁹Department of Environment, Faculty of Environment, Ionian University, 29100 Zakynthos, Greece; ³⁰Dipartimento di Scienze e Tecnologie Biologiche e Ambientali (DiSTeBA), Campus Ecotekne, Strada Provinciale Lecce-Monteroni, 73100 Lecce, Italy; ³¹Institute of Oceanology - Bulgarian Academy of Sciences, Varna, Bulgaria; ³²Conservation Biology Research Group, Department of Biology, University of Malta, Msida, Malta; ³³Institute of Fish Resources - Agricultural Academy, Varna, Bulgaria; ³⁴Faculty of Natural and Agricultural Sciences, Department of Natural Sciences Ovidius University of Constanta, Constanta, Romania; ³⁵Center for Advanced Studies of Blanes (CEAB, CSIC), Carrer Accés Cala Sant Francesc, 14, 17300 Blanes, Girona, Spain; ³⁶Department of Biology, Faculty of Biology, "Alexandru Ioan Cuza" University of Iași, 700507 Iași, Romania; ³⁷Aqua Stories; via Vampolieri, 25/31 int. 27, 95022 Aci Catena (Catania), Italy; ³⁸CNR/ISAFoM (National Research Council of Italy/Institute for Agricultural and Forestry System in the Mediterranean) via Empedocle, 58, 95125 Catania, Italy; ³⁹Frederick University, Limassol, Cyprus; ⁴⁰Department of Integrative Marine Ecology (EMI), Stazione Zoologica Anton Dohrn, Villa Comunale, I-80121 Naples, Italy; ⁴¹NBFC, National Biodiversity Future Centre - Palermo, Piazza Marina 61, I-90133 Palermo, Italy; ⁴²Department of Biology, University of Malta, Msida, Malta; ⁴³Departamento de Botánica y Fisiología Vegetal, Facultad de Ciencias, Campus de Teatinos s/n, 29080 Málaga, Spain; ⁴⁴CMMI - Cyprus Marine & Maritime Institute, 6023, Larnaca, Cyprus; ⁴⁵Department of Biology, Faculty Engineering and Natural Sciences, Manisa Celal Bayar University Muradiye, Manisa 45140, Türkiye; ⁴⁶SAP/RAC, Boulevard du Leader Yasser Arafat, B.P. 337 - 1080 Tunis Cedex - Tunisie; ⁴⁷School of Biological Sciences, University of Aberdeen, Aberdeen AB24 3UU, Scotland, UK; ⁴⁸Institute of Earth Systems, University of Malta, Msida, Malta; ⁴⁹Oceanography Center, University of Cyprus, 1 Panepistimiou Av., 2109 Aglandjia, Nicosia, Cyprus; ⁵⁰Department of Fisheries and Marine Research (DFMR), Ministry of Agriculture, Rural Development and the Environment of the Republic of Cyprus, 101 Vithleem Street, 2033 Strovolos, Nicosia, Cyprus; ⁵¹Departamento de Ecología e Hidrología, Universidad de Murcia, Murcia, Spain; ⁵²Dipartimento di Biologia, Roma Tor Vergata, 00133 Roma, Italy; ⁵³IUCN ROWA, Amman 11194 Jordan; ⁵⁴Mugla Sıtkı Koçman University Faculty of Fisheries, Mugla/Türkiye; ⁵⁵IMC - International Marine Centre, Loc. Sa Mardini, Torregrande - 09170 Oristano, Italy; ⁵⁶Institute for Marine Biological Resources and Biotechnology National Research Council of Italy (IRBIM-CNR) Largo Fiera della Pesca, 2, 60125 Ancona AN, Italy; ⁵⁷Italy Ente Fauna Marina Mediterranea, Avola 96012, Siracusa, Italy; ⁵⁸Department of Earth and Marine Sciences (DiSTeM), University of Palermo, Via Archirafi 20-22, 90123 Palermo, Italy; ⁵⁹Department of Biology, University of Crete, 70013, Heraklion, Greece; ⁶⁰Faculty of Fisheries, Muğla Sıtkı Koçman University, Muğla, Turkey; ⁶¹Department of Biology, Stazione Idrobiologica Umberto D'Ancona, University of Padova, Chioggia, Venice, Italy; ⁶²Department of Biology, University of Padova, Via U. Bassi 58/B, 35121, Padova, Italy; ⁶³Astrolabe-Marine Research, Thessaloniki, 55132, Greece; ⁶⁴Oceanides-Institute of Marine Research & Education, Kalymnos Island, 85200, Greece; ⁶⁵Department of Biology, University of Naples "Federico II", Via Cintia 26, I-80126 Naples, Italy; ⁶⁶Conservation Biology Research Group, Department of Biology, University of Malta, MSD2080 Msida, Malta; ⁶⁷Eser Deniz Ecological, Environmental Company, Technocity of Manisa Celal Bayar University, Muradiye-Manisa, Türkiye; ⁶⁸Institute for Marine Biological Resources and Biotechnologies (IRBIM), National Research Council (CNR), Mazara del Vallo, Italy; ⁶⁹National Institute for Scuba Diving, Zeytouna bay, Beirut; ⁷⁰Arpa Sicilia UOC Area Mare, Lungomare Cristoforo Colombo 4521, Loc. Addaura, 90149 Palermo, Italy; ⁷¹Istanbul University, Faculty of Science, Istanbul/Türkiye; ⁷²Bandirma Onyedil Eylül University, Maritime Vocational School, Balıkesir, Türkiye; ⁷³Department of Biology, Faculty of Marine Sciences, Instituto Universitario de Investigación en Estudios Ambientales y Recursos Naturales IUNAT, Universidad de Las Palmas de Gran Canaria, 35017 Las Palmas, Spain; ⁷⁴Ege University, Faculty of Fisheries, Department of Hydrobiology, 35100 Bornova, İzmir, Türkiye; ⁷⁵Department of Earth and Environmental Sciences, University of Pavia, Italy; ⁷⁶Department Marine Sciences and Applied Biology, University of Alicante, Alicante, Spain; ⁷⁷Department of Biological, Chemical and Pharmaceutical Sciences and Technologies, University of Palermo, 90123 Palermo, Italy; ⁷⁸Dipartimento di Scienze della Terra e del Mare (DiSTeM), Via Archirafi 22, 90123 Palermo, Italy; ⁷⁹Department of Marine Science and Applied Biology, University of Alicante, 03690 Alicante, Spain; ⁸⁰Department of Biosciences, Biotechnologies and Environment, University of Bari Aldo Moro, Via E. Orabona 4, 70125, Bari, Italy; ⁸¹Department of Biology, University of Padova, Via Ugo Bassi 58/B, 35131, Padova, Italy; ⁸²Hellenic Centre for Marine Research, Institute of Marine Biological Resources and Inland Waters, 576 Vouliagmenis Avenue, Argiroupolis, 16452 Athens, Greece; ⁸³Laboratoire de Recherche Ecosystèmes et Ressources Aquatiques, Institut National Agronomique de Tunisie, 43 Avenue Charles Nicolle, 1082 Tunis, Tunisia; ⁸⁴Independent Researcher & International Environmental Consultant, 31400 Toulouse, France; ⁸⁵Department of Science, Roma Tre University, Viale Guglielmo Marconi 446, 00146 Rome, Italy; ⁸⁶Department of Earth, Environment and Life Sciences (DiSTAV), University of Genoa, Corso Europa 26, 16132 Genova, Italy; ⁸⁷Department of Ecology, School of Biology, Aristotle University of Thessaloniki, Thessaloniki, Greece; ⁸⁸Mohammed V University in Rabat, Faculty of Sciences, Laboratory of Biodiversity, Ecology and Genome, 4 Avenue Ibn Battouta, B.P. 1014 RP, Rabat, Morocco; ⁸⁹Natural Sciences and Environment Research Hub, University of Gibraltar, Europa Point Campus, Gibraltar; ⁹⁰Laboratoire Santé, Environnement et Biotechnologie, Faculté des Sciences Ain Chock, Hassan II University of Casablanca, B.P. 5366 Maarif, 20100 Casablanca, Morocco; ⁹¹Arpa Sicilia UOC Area Mare, Lungomare Cristoforo Colombo 4521, Loc. Addaura, 90149 Palermo, Italy; ⁹²Benthic Ecology & Technology

Laboratory, Fisheries Research Institute (ELGO-DIMITRA), 64007, Nea Peramos, Kavala, Greece; ⁹³D.I.I.V. for marine, freshwater and subterranean ecology Sali IV 2, HR-23281 Sali, Croatia; ⁹⁴Department of Biology and Evolution of Marine Organisms, Stazione Zoologica Anton Dohrn (SZN), Naples, Italy; ⁹⁵Institute of Marine Biology, University of Montenegro, Kotor, Montenegro; ⁹⁶University of Zagreb Faculty of Science, Department of Biology, Rooseveltov trg 6, HR-10000 Zagreb, Croatia; ⁹⁷Department of Biology, University of Naples Federico II, Naples, Italy; ⁹⁸Biological Oceanography and Marine Environment Laboratory (LOBEM), Faculty of Biological Science, University of Science and Technology, Houari Boumediene. B.P 32 El Alia 16111 Bab Ezzouar, Algeria; ⁹⁹National Higher School of Marine Sciences and Coastal Management (ENSSMAL), 16320 Algiers, Algeria; ¹⁰⁰Department of Biology and Evolution of Marine Organisms (BEOM), Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Naples, Italy; ¹⁰¹Corfu Scubabots Diving Center, Kontokali Bay hotel, 49100 Corfu, Greece; ¹⁰²Agricultural University of Athens, School of Animal Sciences, Department of Animal Science, Laboratory of Applied Hydrobiology, Iera odos 75, 11855 Athens, Greece; ¹⁰³Department Marine Sciences and Applied Biology, University of Alicante, Alicante, Spain; ¹⁰⁴Aix Marseille Univ, CNRS, IRD, Avignon Univ, IMBE, UMR 7263, Station Marine d'Endoume, Marseille, France; ¹⁰⁵Hellenic Centre for Marine Research, Institute of Marine Biological Resources & Inland Waters, Former US Base at Gournes, P.C. 71500 municipality of Hersonissos, P.O. Box 2214 T.K. 71003 Heraklion Crete, Greece; ¹⁰⁶Department of Life and Environmental Sciences, Polytechnic University of Marche, Ancona, Italy; ¹⁰⁷Stazione Zoologica Anton Dohrn, Naples, Italy; ¹⁰⁸Institute of Biomolecular Chemistry, National Research Council of Italy, Pozzuoli, Naples 80078, Italy; ¹⁰⁹ISEM, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France; ¹¹⁰Department of Research Infrastructures for Marine Biological Resources, Stazione Zoologica Anton Dohrn, Napoli, Italy; ¹¹¹Institut Océanographique Paul Ricard, Ile des Embiez, 83140 Six-Fours-les-Plages, France / and Université de Toulon, Aix Marseille Univ, CNRS, IRD, MIO, 83130 La Garde, France; ¹¹²Université de Toulon, Aix Marseille Univ, CNRS, IRD, MIO, 83130 La Garde, France; ¹¹³Museo del Mar, Ceuta, Spain; ¹¹⁴Maison de la mer - Aire Marine Protégée de la côte agathoise, Rue de la Capitainerie, 34300 Cap d'Agde – France; ¹¹⁵Animate your Science, Adelaide, Australia; ¹¹⁶Lebanon project office (IUCN, ROWA), Beirut, Lebanon; ¹¹⁷Oceana, Spain; ¹¹⁸Biology Department, The American University in Cairo, P.O. Box 74, New Cairo 11835, Egypt; ¹¹⁹Cretaquarium, Hellenic Center for Marine Research, P.O. Box 2214, 71003 Heraklion, Crete, Greece; ¹²⁰Department of Ecosustainable Marine Biotechnology, Stazione Zoologica Anton Dohrn, Via Ammiraglio Ferdinando Acton 55, 80133 Naples, Italy; ¹²¹Sorbonne Université, CNRS, UMR 1744, Station Biologique, Roscoff, France; ¹²²Aix Marseille Univ, CNRS, IRD, Avignon Univ, TELEMME, UMR 7263, Station Marine d'Endoume, Marseille, France; ¹²³Institute for Marine Biological Resources and Biotechnology (IRBIM), National Research Council – CNR, Mazara del Vallo, Italy; ¹²⁴Smithsonian Environmental Research Center, Edgewater, MD, USA; ¹²⁵Le Corbusier 644, 280 Boulevard Michelet, 13008 Marseille, France; ¹²⁶Mohammed V University in Rabat, Faculty of Sciences, Laboratory of Biodiversity, Ecology and Genome, 4 Avenue Ibn Battouta, B.P. 1014 RP, Rabat, Morocco; ¹²⁷Aix Marseille Univ, CNRS, TELEMME, UMR 7303, Aix-en-Provence, France; ¹²⁸Portofino Marine Protected Area, Viale Rainusso 1, 16038 Santa Margherita Ligure, Italy; ¹²⁹Apulian Regional Agency for the Environmental Prevention and Protection, 70126, Bari, Italy; ¹³⁰Dipartimento di Scienze, Università 'Roma Tre', Viale G. Marconi 446, I-00146 Rome, Italy; ¹³¹Department of Fisheries and Marine Research (DFMR), Ministry of Agriculture, Rural Development and the Environment of the Republic of Cyprus, 2033, Nicosia, Cyprus; ¹³²Department of Integrative Marine Ecology (EMI), Stazione Zoologica Anton Dohrn, Ischia Marine Centre, Ischia, Naples, 80077, Italy; ¹³³Institute of Oceanology - Bulgarian Academy of Sciences, 40 Parvi mai str, 9000 Varna, Bulgaria; ¹³⁴Marine Biology Museum "Pietro Parenzan", Via A. Vespucci, 13/17, 73010 Porto Cesareo (LE), Italia.

Corresponding author: Jennifer C.A. Pistevo (jennie@aegean.gr)

Citation: Pistevo JCA, Zenetos A, Stranga Y, Teaca A, Begun T, Trkov D, Langeneck J, Muresan M, Comas-González R, Sempere-Valverde J, Toso A, Chebaane S, Navarro-Barranco C, Rosso A, Png-Gonzalez L, Giakoumi S, Vázquez-Luis M, Cañizares-González JM, Siddiolo C, Gemma Donato, Giacobbe S, Carbonell A, Zakhama-Sraieb R, Di Martino E, Ammar I, Cosentino A, Mnasri Affi I, Sanfilippo R, Bitar G, Kytinou E, Tari G, Huseyinoglu MF, Gerakaris V, Issaris Y, Lardi P, Oikonomidis G, Salomidi M, Tsirintanis K, Kleitou P, Constantinou G, Michail C, Bariche M, Deidun A, Marrone A, Ragkousis M, Falautano M, Castriota L, Morri C, Bianchi CN, Skouradakis G, Nash FT, Digenis M, Mancini E, Stefanova ES, Vella A, Vella N, Mihneva VV, Papadimitriou E, Skolka M, Stefanova K, Fortić A, Garcia M, Dailianis T, Tamburello L, Surugiu V, Stancanelli B, Di Martino V, Bosmali A, Mavrić B, Gerovasilieou V, Kletou D, Crocetta F, Rallis I, Evans J, Jeschke MA, Cai LL, Taşkın E, Franzitta G, Sghaier YR, Kennedy A, Bartolo AG, Küpper FC, Rousou M, Orenes-Salazar V, Bonifazi A, Hadjioannou L, Vlantaridis Z, El Shaer H, Çiftçioglu M, Cerim H, Lipej L, Grech D, Zotou M, Azzuro E, Schembri PJ, Tiralongo F, Chatzigeorgiou G, Di Lorenzo M, Vigo Fernandez M, Di Franco A, Orlando-Bonaca M, Calò A, Malefaki C, Garcia-Charton JA, Papadakis O, Yapici S, Tanduo V, D'Annello I, Munari M,

Abstract

To enhance regional knowledge of invasive alien species (IAS), cryptogenic, and neonative species, this study compiles previously unpublished records from the Mediterranean and Black Seas via contributions from 227 marine experts. The open-access dataset includes 7,422 records from 21 countries, representing 275 taxa: 222 Animalia, 44 Plantae, and 9 Chromista. Seventy-two percent of records were collected in 2021–2024 while most reports were from Lebanon, Greece, Cyprus, Italy, and Romania. The most frequently recorded species were *Siganus luridus*, *S. rivulatus*, *Callinectes sapidus*, *Caulerpa cylindracea* and *Parupeneus forsskali*. Several records are biogeographically notable: basin-level firsts include *Dynoides amblysinus* and *Symplegma rubra*, and the first confirmed occurrence of *Malmgrenia mcintoshii*. National firsts include *Mesanthura cf. romulea*, *Mitrella psilla*, and *Syllis ergeni* (Algeria); *Exogone brevirantennata*, *Jassa marmorata*, *Leodice antennata*, and *Syllis schulzi* (Cyprus); *Botrylloides niger* and *Polyandrocarpa zorritensis* (Greece); *Microcosmus exasperatus*, *Phallusia nigra*, and *Stypopodium schimperi* (Italy); *Chaetozona corona* (Slovenia); *Pseudodiaptomus marinus* and *Symplegma brakenhielmi* (Spain); *Aquilonastra burtoni*, *Siphonaria crenata*, and *Sphaeroma walkeri* (Syria) and *Distaplia bermudensis* (Tunisia). In the Black Sea, *Hydroides dianthus*, *Ferosagitta* sp., and a zoea III of *Palaemon macrodactylus* are newly documented from Romanian waters. Beyond listing records, this compilation supports spatiotemporal analyses relevant to management. The sharp post-2021 increase reflects both species spread and intensified surveys, and habitat-labelled records include harbours/marinas and shellfish aquaculture settings, consistent with pathway hotspots. By revealing where and when detections concentrate—and which taxa dominate—these data can guide early detection, risk prioritization, and targeted biosecurity, while identifying under-reported coastlines.

Key words: non-native species, marine bioinvasions, inventories, distribution, geo-referenced records, Mediterranean Sea

Kampouris TE, Putignano M, Virgili R, Agius Darmanin S, Bilgiç F, Sardo G, Calabrò M, Nikolaou A, Erga Z, Zogheib A, Balistreri P, Çağlar S, Yalçın F, Osca D, Gratsia E, Aglieri G, Çinar ME, Motalipassi M, Coco S, Gkoulia A, Ramos-Esplà AA, Mannino AM, Dimichele D, Spinelli S, Arroyo Martínez E, Tursi A, Capasso A, Christidis G, Solca M, Rifi M, El Zrelli RB, Bouhlef R, Scalera R, Bandi A, Reboa A, Mazaris A, Bazairi H, Bertolino M, Selfati M, Sirchia B, Georgiadis CG, Petricoli D, Macaluso E, D'Aniello E, Fotiadis N, Tezel R, Petović S, Bakran-Petricoli T, Pitacco V, Colletti A, Bensari B, Bahbah L, D'Aniello S, Psailas S, Kalogirou S, Kaikai T, Forcada A, Manfredonia A, Bottalico A, Valle C, Bedotto C, Damalas D, Sacco D, Mollo E, Iatrou F, Cerqueira F, Viard F, Villani G, Foulquié M, Geraci ML, Šiško M, Ocaña O, Dupuy de la Grandrive R, Vitale S, Rossi T, Samaha Z, Aguilar R, Alvarez H, Menabit S, Pavel AB, Azzola A, Bos AR, Anastasiadis A, Blanco J, Lejeune C, Coppola D, Davoult D, Faget D, Scannella D, Keppel E, Falsone F, Palma Esposito F, García S, Furfaro G, Zibrowius H, Iziani I, Anastasopoulou I, Kavakakis K, Scavino L, Merotto L, Yankova M, Montefalcone M, Leban N, Ungaro N, Mariottini P, Perry AL, Stavrou P, Mazzella V, Raykov V, Ferrario J, Stamouli C, Siapatis A, Vernadou E, Bommarito C, Milisenda G, Somma E, Russo S, Katsanevakis S (2026) Unpublished Mediterranean and Black Sea records of marine alien, cryptogenic, and neonative species (III). *BioInvasions Records* 15(3): 533–572, <https://doi.org/10.3391/bir.2026.15.3.01>

Received: 9 September 2025

Accepted: 20 March 2026

Published: 25 May 2026

Thematic editor: Andrew Davinack

Copyright: © Pistevo et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International - CC BY 4.0).

OPEN ACCESS

Introduction

Biological invasions represent one of the most significant and fast-growing threats to marine biodiversity worldwide, often leading to the disruption of native ecosystems and ecosystem services (Molnar et al. 2008; Katsanevakis et al. 2014; Tsirintanis et al. 2022). The Mediterranean Sea, despite its relatively small size, has emerged as a hotspot for marine invasive alien species (IAS) and hosts the highest number of introductions among European seas (Galil et al. 2014; Thomsen et al. 2016; Seebens et al. 2016; Zenetos et al. 2022a). This phenomenon is facilitated by a combination of intense anthropogenic pressures, including maritime traffic, aquaculture, and the Suez Canal, as well as regional warming that enhances the suitability of conditions for thermophilic species (Zenetos et al. 2012; Schroeder et al. 2016; Albano et al. 2021). Due to its geopolitical complexity, being bordered by 22 countries with varying policies and capacities for marine monitoring, the Mediterranean faces significant challenges in implementing cohesive biosecurity and biodiversity monitoring frameworks (Zenetos and Galanidi 2020). Meanwhile, the neighbouring Black Sea has experienced a steady rise in alien species introductions over the past five decades, driven both by the secondary spread of IAS from the Mediterranean via the Dardanelles and the Bosphorus, and by primary introductions, mainly through shipping (Băncilă et al. 2022; Ragkousis et al. 2023).

Increasing attention is also being paid to neonative species—those undergoing climate-driven natural range expansions—which can also exert ecological pressures (Essl et al. 2019; Azzurro et al. 2022; Virgili et al. 2025). Whether alien or neonative, their expanding presence continues to drive processes such as tropicalization in the Mediterranean (Bianchi and Morri 2003; Peleg et al. 2020) and Mediterraneanization in the Black Sea (Kideys et al. 2000; Eyuboglu 2022).

Despite the ecological significance of these changes, a large volume of valuable distribution data remains unpublished or inaccessible, often stored in personal archives or institutional databases (Katsanevakis et al. 2020). This gap hinders comprehensive assessments of invasion dynamics, species trends, and the development of management responses under frameworks such as the European Marine Strategy Framework Directive (MSFD) and the Barcelona Convention/UNEP/MAP (Tsiamis et al. 2019; 2021). The need for frequent updates to alien species inventories has led to coordinated efforts to collect and disseminate such data via collaborative networks of marine scientists.

This paper builds upon two previous collaborative efforts to document unpublished records of alien species – including IAS in the region. The first effort compiled 5,376 records of 239 alien and cryptogenic marine taxa from the Mediterranean Sea, contributed by 126 scientists across 16 countries (Katsanevakis et al. 2020). A subsequent expansion incorporated

both the Mediterranean and Black Seas, collecting 12,649 records of 247 taxa, including neonative species, and substantially enriching the spatial and temporal resolution of available data (Ragkousis et al. 2023). The present study represents the third iteration of this initiative and aims to further strengthen regional inventories by collating newly gathered and previously unpublished records of NIS, cryptogenic, and neonative species. In doing so, it contributes to a growing open-access dataset that is essential for tracking bio-invasion dynamics and supporting efforts to monitor, manage, and mitigate the impacts of alien species in the region.

Dataset compilation

Following the approach of the previous two iterations of this initiative, the call for contributions was circulated by direct e-mail invitation to known marine experts across Mediterranean and Black Sea countries, using a standardized Excel template to harmonize submissions. Compared to the template used in our previous studies, we revised several terms to better align with the Darwin Core standards (<http://rs.tdwg.org/dwc/doc/em/>). Contributors were also invited to suggest additional expert contacts, enabling further targeted outreach through professional networks. Records originating from third parties (including citizen-science platforms) were included only when the identification was validated by a marine expert, typically supported by photographic documentation and/or reliable observer provenance as reported in Supplementary material Table S1. No records derived from fisher questionnaires were included in the present compilation.

Overall, a total of 227 contributors from 21 different countries provided data, resulting in a dataset of 7,422 individual records (see Table S1). Each entry in the spreadsheet corresponds to a unique observation, documented with spatial and temporal details. Key information included species name, biogeographic status, geographic coordinates (latitude and longitude), country of observation, year, name of the identifier, observation type, and documentation method.

Biogeographic status followed established references—Zenetos et al. (2022b) for the Mediterranean and Băncilă et al. (2022) for the Black Sea—supplemented by species-specific literature when classification was uncertain. Four categories were used to define status: “Alien” (as per Essl et al. 2018), “Neonative” (*sensu* Essl et al. 2019), “Cryptogenic” (unknown origin; Carlton 1996; Essl et al. 2018), and “Data deficient” (for taxa whose biogeographic status could not be determined due to lack of data; Essl et al. 2018). Neonatives were included as a distinct status category to capture climate-driven, natural range expansions (*sensu* Essl et al. 2019), and to avoid conflating such range recognition with human-mediated introductions. In the Mediterranean and Black Sea, neonatives may expand through natural corridors (e.g., through the Gibraltar Strait or into the Black Sea via

the Dardanelles–Sea of Marmara system), as discussed in the previous iteration of this initiative (Ragkousis et al. 2023). However, we do not attempt to reconstruct entry routes or quantify range-shift trajectories for each neonative taxon, as this would require standardized baseline distributions and longitudinal time-series beyond the scope of this record-mobilization dataset.

Record verification and inclusion criteria

The dataset comprises records submitted by participating marine experts (227 contributors from 21 countries). Only records with the required minimum information were retained (taxon name, biogeographic status, coordinates/country, year, identifier, observation type, and documentation method). “Observation type” describes the verification pathway and includes (i) direct expert visual identification (e.g., in situ surveys, including validated photographic material submitted to the expert by collaborators) and (ii) specimen-based identification following collection by scientific personnel, typically confirmed through morphological examination in the laboratory and, where necessary/available, DNA-based confirmation.

Optional fields were used when provided to capture additional context (e.g., specific or estimated observation date, depth, habitat type, estimated abundance or coverage, and supporting notes). The dataset is dominated by conspicuous and readily identifiable alien taxa; for biogeographically notable records (e.g., basin-level or national firsts) and/or taxa with higher identification uncertainty, supporting details are provided within the relevant species accounts. Where the available evidence was insufficient to confidently resolve taxonomic identity and/or biogeographic status, records were treated conservatively using the dataset categories (e.g., cryptogenic).

Taxonomic coverage

A total of 275 distinct taxa were recorded in the dataset, primarily Animalia (222; 80.58%), with smaller proportions of Plantae (44; 15.98%) and Chromista (9; 3.44%) (Figure 1). Within Animalia, records were dominated by Chordata (primarily fishes), followed by major invertebrate groups (notably Arthropoda and Mollusca) (Figure 1C). This dominance of conspicuous taxa—especially fishes—is consistent with previous iterations of this initiative and likely reflects both detectability and monitoring focus. Most of the records were classified as alien (6,438; 86.6%), with additional contributions from cryptogenic (752; 10.1%), neonative (84; 1.1%), and uncertain cases assigned as neonative/cryptogenic (161; 2.2%). The low proportion of neonative records (1.1%) should not be interpreted as evidence that climate-driven range expansions are rare in the region. Rather, it likely reflects the purpose and reporting bias of this expert-mobilized compilation, which was primarily designed to strengthen alien species inventories (the




Figure 1. Summary of taxonomic information from the dataset: (A) proportion of records grouped by kingdom; (B) histogram showing the distribution of record counts per species; (C) breakdown of records by phylum; (D) species-level record frequency for the 20 taxa with the highest number of observations.

first iteration of this initiative targeted alien and cryptogenic species specifically) and therefore tends to elicit and prioritize reports of alien taxa. In addition, identifying a species as neonative requires stronger contextual information (baseline distribution knowledge and confidence that the expansion is natural rather than human-mediated), which increases uncertainty and can lead to conservative categorization. This is reflected in the additional 2.2% of records classified as “neonative/cryptogenic”, i.e., cases where species status is uncertain.

Approximately 77% of the reported taxa come from five countries—Lebanon, Greece, Cyprus, Italy, and Romania. The five species with the highest number of records were *Siganus luridus* (Rüppell, 1829) (447), *Siganus rivulatus* Forsskal & Niebuhr, 1775 (435), *Callinectes sapidus* Rathbun, 1896 (303), *Caulerpa cylindracea* Sonder (269), and *Parupeneus forsskali* Fourmanoir & Guézé, 1976 (263).

Spatial and temporal coverage

The temporal span of the dataset ranges from 1973 to 2025. A substantial majority of records (approximately 72%) were collected between 2021 and 2024, with the highest contributions in 2024 (1,859 records), followed by 2023 (1,665), and 2022 (1,286) (Figure 2A). The dataset includes records from 21 countries, including 6,384 records from the Mediterranean Sea and 1,046 from the Black Sea (Figure 2B). The species recorded in the highest number of countries were *Callinectes sapidus* Rathbun, 1896 (10), *Bursatella leachii* Blainville, 1817 (8), *Fistularia commersonii* Rüppell, 1838 (8), *Amathia verticillata* (delle Chiaje, 1822) (7), *Caulerpa cylindracea* Sonder (7), *Pinctada radiata* (Leach, 1814) (7), and *Pterois miles* (Bennett, 1828) (7) (Figure 2C). Establishment status was not evaluated systematically in this study because it is scale-dependent and not uniquely attributable at the record level; readers are referred to the most recent basin-level assessments for the Mediterranean and Black Sea (Zenetos et al. 2022b; Magliozzi et al. 2025).

The spatial distribution of records remains uneven across the region (Figure 3A), influenced by species occurrence patterns, variability in sampling effort, and contributor distribution. In addition to uneven sampling effort and data-sharing, spatial gaps can also reflect uneven availability of taxonomic expertise among countries and taxa; absence of records should therefore not be interpreted as evidence of absence. A substantial portion of the older unpublished records in our dataset originated from Lebanon, which also accounts for the highest overall number of records (Figure 2B). The inclusion of historical observations (1970s–1990s) does not reflect a timeline inconsistency, but rather the purpose of this initiative to mobilize and georeference information that is often unpublished or inaccessible (e.g., retained in personal archives or institutional databases). As noted in previous iterations, alien-species observations are often published only when they constitute “first records”, whereas many subsequent detections (and older observations) remain scattered and unpublished, despite their value for reconstructing invasion trajectories and temporal dynamics. In addition, some historical findings—particularly from Lebanon—have been mentioned previously without sufficient spatial detail for reuse; the present compilation provides the missing georeferenced locality information where available and includes backdated historical records not previously published (e.g., the three macroalgae reported as backdated first country records for Lebanon).

High reporting volumes were also observed in Greece, Cyprus, Italy, and Romania. Conversely, record contributions were scarce in Albania, Egypt, Georgia, Ukraine, France, and parts of North Africa, with only isolated records submitted from these regions (Figure 3A). This pattern likely reflects differences in sampling effort and data sharing rather than the actual absence or low presence of alien and other species targeted in the present study. The combination of the mapping points originating from all the three studies




Figure 2. Spatial and temporal distribution characteristics of the NIS dataset: (A) yearly distribution of recorded observations; (B) number of records contributed by each country; (C) number of countries reporting each species (shown for species recorded in five or more countries).




Figure 3. (A) number of alien, cryptogenic, and neontive records per 10 km × 10 km grid cell, included in the dataset of this study; (B) cumulative number of alien, cryptogenic and neontive records with the same grid cell from this study plus Katsanevakis et al. (2000) and Ragkousis et al. (2023).

(Katsanevakis et al. 2020; Ragkousis et al. 2023; present data) highlights both the extent of species spread and the areas where fewer records are evident (Figure 3B), which may reflect either information gaps or genuinely lower presence of IAS (e.g., parts of Sardinia and the Tyrrhenian coastline).

Remarkable new records of alien species

This dataset includes several records of particular significance, including first basin-level, national, and sub-regional occurrences, as well as confirmations of species establishments. Notably, the isopod *Dynoides amblysinus* Pillai, 1954, is recorded for the first time in the Mediterranean Sea, based on material collected from the Port of Catania (Italy). Other basin-level firsts include the colonial ascidian *Symplegma rubra* Monniot, 1972, newly observed in the Mediterranean (off the coast of Tunisia), and the polychaete *Malmgrenia mcintoshi* (Tebble & Chambers, 1982), which is here reported for the first time from the Ionian Sea, confirming its presence in the central Mediterranean Basin. *Paracalanus tropicus* Andronov, 1977 is recorded for the first time in Mediterranean waters in Algeciras Harbour, Spain. *Stypopodium schimperi* (Kützing) Verlaque & Boudouresque, a brown alga previously limited to the eastern Mediterranean, is reported for the first time in southern Italy, specifically from Lampedusa and Pantelleria in the Sicily Strait.

Several records represent first observations at the national level. From Syria, the starfish *Aquilonastra burtoni* (Gray, 1840), the limpet *Siphonaria crenata* Blainville, 1827, and the isopod *Sphaeroma walkeri* Stebbing, 1905 are reported for the first time. From Cyprus, new records include the polychaetes *Exogone brevantennata* Hartmann-Schröder, 1959, *Leodice antennata* Savigny in Lamarck, 1818 and *Syllis schulzi* (Hartmann-Schröder, 1960), and the amphipod *Jassa marmorata* Holmes, 1905. In Algeria, new findings include the polychaete *Syllis ergeni* Çinar, 2005, the isopod *Mesanthura* cf. *romulea* Poore & Lew Ton, 1993, and the gastropod *Mitrella psilla* (Duclos, 1840). The cirratulid polychaete *Chaetozone corona* Berkeley & Berkeley, 1941 is reported for the first time from Slovenian waters. In Tunisia, first records include the colonial ascidian *Distaplia bermudensis* Van Name, 1902, and an occurrence of *Polyandrocarpa zorritensis* Van Name, 1931 from 2021, which backdates the first published Tunisian record by three years (Melki et al. 2026). This backdating is consistent with the generally short reporting lags documented for Tunisia and the comparatively longer (often near-decadal) lags reported for ascidians (Zenetos et al. 2019; Galanidi et al. 2025). In Italy, new national or remarkable records regard the confirmation of the ascidians: *Phallusia nigra* Savigny, 1816, which was only known in DNA samples (Varella et al. 2025), *Microcosmus exasperatus* Heller, 1878, *Distaplia bermudensis* Van Name, 1902, and *Polyclinum constellatum* Savigny, 1816; the mollusc *Lamprohaminoea ovalis* (Pease, 1868); and the polychaete *Eurythoe complanata* (Pallas, 1766),

the latter four recorded in the Adriatic Sea for the first time. The ascidians *Botrylloides niger* Herdman, 1886 and *Polyandrocarpa zorritensis* (Van Name, 1931) are documented for the first time in Greece. In Spanish Mediterranean waters, the tunicate *Symplegma brakenhielmi* (Michaelsen, 1904) is here reported from Alicante port while the copepod *Pseudodiaptomus marinus* Sato, 1913 is recorded for the first time from the Algeciras Harbour, representing the westernmost known occurrence of the species in the basin.

In the Black Sea, the present dataset provides the first records for Romania of the third-stage larvae of *Palaemon macrodactylus* Rathbun, 1902, the serpulid polychaete *Hydroides dianthus* (Verrill, 1873), and a new chaetognath of the genus *Ferosagitta*. It also provides further evidence of the establishment of the hydrozoan *Podocorynoides minima* (Trinci, 1903) in Romanian coastal and offshore habitats (Mureşan et al. 2021).

Additional significant range extensions include the bryozoans *Catenicella paradoxa* Rosso, 2009 and *Tricellaria inopinata* d'Hondt & Occhipinti-Ambrogi, 1985, the former first reported from the Sicily Strait and the latter observed for the first time in Sicily and the western Ionian Sea. The bryozoan *Arbopercula angulata* (Levinsen, 1909) is newly reported from the harbour of Lerici (Liguria, Italy), representing the northernmost known occurrence in both the Tyrrhenian and western Mediterranean. The spionid polychaete *Prionospio depauperata* Imajima, 1990 is also newly recorded from the Tyrrhenian Sea (western Mediterranean), extending its range beyond its previously known distribution in the eastern and central Mediterranean (Dağlı and Çinar 2009; Ragkousis et al. 2023; Langeneck et al. 2024).

Additionally, three macroalgae and one fish are here reported as backdated first country records for Lebanon: *Codium taylorii* P.C. Silva, *Galaxaura rugosa* (J.Ellis & Solander) J.V. Lamouroux, *Padina boergesenii* Allender & Kraft, and *Terapon puta* Cuvier, 1829, based on historical records not previously published.

A recent study of the structure of marine biofouling in two fishing and recreational ports in southern Syria has revealed more established alien species. In this report, we record the presence of the starfish *Aquilonastra burtoni* (Gray, 1840), the limpet *Siphonaria crenata* Blainville, 1827, and the isopod *Sphaeroma walkeri* Stebbing, 1905 for the first time in the country.

In the following taxon-focused subsections, we briefly present selected records of particular biogeographic importance (e.g., first records for the Mediterranean/Black Sea and/or first country records) and provide the available documentation supporting their identification, following the approach used in previous iterations of this initiative.

Polychaeta (Annelida)

- *Malmgrenia mcintoshii* (Tebble & Chambers, 1982) – first record in the Ionian Sea (central-eastern Mediterranean).

A single adult specimen of the polychaete *M. mcintoshii* has been recorded in July 2006 in shallow waters (2 m depth) in the Catania harbour area (Sicily, Italy). The specimen was found settled inside a living tube of *Chaetopterus variopedatus* (Renier, 1804), on a mixed sandy-rocky bottom, affected by temporary accumulation of pyroclastic sands. Identification was confirmed by Dr. Ruth Barnich. The specimen is stored at the Senckenberg Museum Polychaete Collection (code ID sesam-248664, catalogue no. 16988), preserved in 70% ethanol.

Malmgrenia mcintoshii is generally regarded as an Atlantic–Boreal species, with a main distribution in the northeastern Atlantic extending to the Barents Sea (Marin and Antokhina 2022). In a biogeographical revision of the genus *Malmgrenia*, its presence was generically stated for the northeastern Atlantic and the Mediterranean Sea (Barnich et al. 2017), whereas Marin and Antokhina (2022) considered it currently absent from the Mediterranean.

The present record therefore represents the first documented occurrence for Italian waters and confirms presence in the Mediterranean at least in the Ionian Sea (central-eastern basin). The apparent uncertainty in its Mediterranean distribution likely reflects its symbiotic lifestyle: species of *Malmgrenia* are commensals within tubicolous macrofauna (Martin and Britayev 1998; cited here for ecological context), and *M. mcintoshii* is typically associated with large *Chaetopterus* tubes. Such ecological specificity can lead to punctiform and irregular occurrences and may contribute to historical under-detection or misidentification. At the same time, its presence in a harbour setting with moderate anthropogenic disturbance means that passive transport (e.g., of juvenile stages) cannot be excluded. Accordingly, its status in the Mediterranean is best treated as cryptogenic rather than alien.

- *Exogone breviantennata* Hartmann-Schröder, 1959 – first record in Cyprus.

The polychaete *E. breviantennata* was first reported in the Mediterranean Sea from shallow-water vegetated environments off Lebanon (Aguado and San Martín 2007) and subsequently from Türkiye (Çinar 2009). This species is morphologically very similar to the native *Exogone verugera* (Claparède, 1868) and the alien *Exogone africana* Hartmann-Schröder, 1974. It can be distinguished from *E. verugera* by the presence of a dorsal cirrus at the second chaetiger, a character shared with *E. africana*, which however has a longer proventricle. The occurrence off Cyprus represents the first record of *E. breviantennata* for this island.

- *Leodice antennata* Savigny in Lamarck, 1818 – first record in Cyprus.

The polychaete *L. antennata* (Figure S1) is a Lessepsian species currently known with certainty for Türkiye, Egypt, and Lebanon (Kurt Şahin and Çinar 2009, 2017; Dorgham and Hamdy 2015; Fortič et al. 2023). The recorded specimens represent the first report from Cyprus. A COI

sequence obtained from one specimen (GenBank accession code: PV018846) has a 99% identity with a specimen from Lebanon sequenced by Toso et al. (2024) (Figure S1).

- *Syllis ergeni* Çinar, 2005 – first record in Algeria.

The polychaete *S. ergeni* (Figure S2) is a syllid species often associated with polluted and anthropized environments in the eastern Mediterranean Sea (Çinar et al. 2008, 2017; Rousou et al. 2023). Even if it was originally described from the Mediterranean Sea (Çinar 2005), it was subsequently recorded in the Red Sea and elsewhere in the Indo-Pacific region (Çinar et al. 2021; San Martín et al. 2024) and consequently considered alien in the Mediterranean basin (Çinar et al. 2017). The present records from Algeria considerably extend its distribution towards the western Mediterranean Sea.

- *Syllis schulzi* (Hartmann-Schröder, 1960) – first record in Cyprus.

The polychaete *S. schulzi* (Figure S3) was described for the Red Sea and later reported in the western Mediterranean Sea. Zenetos et al. (2010) and Çinar (2013) considered it as a questionable alien species. The distribution of *S. schulzi* encompasses the southern Mediterranean Sea as well as the Red Sea (Ba-Akdah et al. 2018; Toso et al. 2024), suggesting that it should be considered cryptogenic in the basin. It is here recorded for the first time from the coast of Cyprus.

- *Prionospio depauperata* Imajima, 1990 – westwards expansion.

Prionospio depauperata Imajima, 1990, (Figure S4) is a spionid polychaete originally described from Japan and widely reported from the eastern and central Mediterranean Sea (Dağlı and Çinar 2009; Ragkousis et al. 2023; Langeneck et al. 2024). The present data extend its distribution to the Tyrrhenian Sea, western Mediterranean.

- *Chaetozone corona* Berkeley & Berkeley, 1941 – first record in Slovenia.

Chaetozone corona (Figure S5) is a bitentaculate cirratulid polychaete that is considered an established alien species along the coasts of Europe (Le Garrec et al. 2017; Langeneck et al. 2020). The species can be easily distinguished from its congeners owing to its spines starting from the first chaetigers. It was originally reported from the Pacific Ocean (Berkeley and Berkeley 1941) and is currently known from several areas of the Mediterranean from where it was first recorded in muddy sands in the Bay of Izmir (Çinar and Ergen 2007), and later in the eastern coast of the Adriatic Sea (Grossi et al. 2017; Munari et al. 2017). This species was probably introduced from its original range via shipping (Le Garrec et al. 2017). In Slovenia, specimens of *C. corona* were found during benthic surveys as part of the national monitoring programme at four sites in the southern part of the Gulf of Trieste (Slovenia, northern Adriatic Sea) using

a Van Veen grab (0.1 m²). The first specimen was found on June 2, 2022, on soft bottom at a depth of 7.5 m and three more specimens in 2023: one on June 2 at a depth of 7.5 m and the others at two sites at a depth of 8.1 m and 8.3 m.

- *Hydroides dianthus* (Verrill, 1873) – first record in Romania.

The serpulid polychaete *H. dianthus* is known for its role as a biofouler and invasive species across a wide range of marine and brackish environments globally (Sun et al. 2017). Initially described from the U.S. East Coast, its distribution has expanded significantly, now including the Mediterranean, Black Sea, Pacific, and Indian Oceans (Zibrowius 1973; Bastida-Zavala and ten Hove 2002; Streftaris et al. 2005), recently extending to northern China and Japan (Sun and Yang 2000; Link et al. 2009). In the Black Sea, it was first recorded along the Sevastopol coast, Crimea (Boltacheva et al. 2011); and later from the Sea of Marmara by Çinar et al. (2020, 2024). Here, we report the first documented occurrence of *H. dianthus* along the Romanian Black Sea coast, collected in September 2024 at a depth of 7 m on sandy-mud substrates (Figure S6). Morphological features of the specimen matched the original species description. The operculum exhibits a distinctive two-tiered structure functioning as a plug for the calcareous tube, comprising a basal funnel and a distal spinous verticil (Figure S6B). The funnel features 27 radii with pointed tips. Collar chaetae bear two short blunt teeth and a smooth distal blade. The tube is marked by two longitudinal ridges and lacks peristomes or transverse ridges. The branchial crown includes approximately 16 radioles on each side of the mouth, forming about one-quarter of the worm's total length.

The site of this finding in proximity to the Port of Constanța suggests introduction via maritime traffic. This aligns with recent first Romanian records of other alien polychaetes, such as *Laonome xeprovala* Bick & Bastrop, in Bick et al. 2018 and *Marenzelleria neglecta* Sikorski & Bick, 2004 (Teacă et al. 2021, 2022; Ragkousis et al. 2023), which are also likely shipping-related introductions. The ongoing spread of *H. dianthus* emphasizes the need for constant monitoring, research into its biology and invasion pathways, and the development of effective, environmentally sustainable control measures.

Crustacea (Isopoda, Copepoda and Decapoda)

- *Dynoides amblysinus* Pillai, 1954 –first record in the Mediterranean Sea (Port of Catania, Italy).

The isopod crustacean genus *Dynoides* Barnard, 1914 currently includes 19 species distributed across tropical and subtropical regions (Boyko et al. 2025; Espinosa-Pérez and Hendrickx 2002; Khalaji-Pirbalouty and Gagnon 2021). *Dynoides amblysinus* is native to tropical and temperate seas and has been reported from the western Indian Ocean to eastern Australia, Hong

Kong, and Japan (Bruce 1982; Harrison and Holdich 1984; Li 2000; Nunomura 2010; Kazmi and Yousouf 2013; Bruce and Wong 2015). In July 2024, during the vagile fauna sampling from two PVC panels, *D. amblysinus* was recorded at a depth of 1.2 m on the quay wall of the Levante Pier in the Port of Catania (Figure S7). This finding represents the first known record of the species in the Mediterranean Sea.

No specific information is currently available on the ecology of *D. amblysinus* but studies on other species of the same genus provide some general insights. For example, *D. crenulatus* Carvacho & Haasmann, 1984 and *D. saldanai* Carvacho & Haasmann, 1984, found along the Pacific coast of Mexico, inhabit rocky bottoms of lower intertidal zones thriving among algae such as *Jania adhaerens* J.V. Lamouroux; *Amphiroa misakiensis* Yendo; *Chaetomorpha linum* (O.F.Müller) Kützing and *Hypnea pannosa* J. Agardh (Espinosa-Pérez and Hendrickx 2002).

- *Sphaeroma walkeri* Stebbing, 1905 – first record in Syria.

Sphaeroma walkeri (Figure S8) is a globally distributed isopod (WoRMS Editorial Board 2025), native to the Indian Ocean and the Red Sea. It is a widely ship-transported alien that entered the Mediterranean Sea as far back as 1924 (Omer-Cooper 1927). It is found on many Mediterranean coasts, including the eastern Mediterranean (e.g., the Aegean Sea, Greece) and Tunisia (Galil 2008; Molnar et al. 2008; Ounifi-Ben Amor et al. 2015; Zenetos et al. 2020). It thrives in crevices, under stones and wood, and is common in fouling communities. Many specimens were found among individuals of the invasive mussel *Brachidontes pharaonis* (P. Fischer, 1870) on the Latakia coast at the sites of Afamia, near the fishing port, and the southern Kornish, accompanied by many polychaetes. Small individuals, less than 5 mm long, were observed in densities of 10–30 individuals per m² from March to August 2021 in the intertidal zone. The body bears distinct tubercles on the telson and pods, and the telson is toothed (Figure S8A–C). This isopod appears to have been established on the Syrian coast long ago, but its discovery was delayed, probably due to its cryptic habit.

- *Mesanthura cf. romulea* Poore & Lew Ton, 1993 – first record in Algeria.

The cryptogenic *Mesanthura cf. romulea* (Figure S9) is currently widespread in the Mediterranean Sea, where it locally reaches considerable abundances (Ulman et al. 2017). While already known for most Mediterranean countries, it is here reported for the first time from Algeria.

- *Palaemon macrodactylus* Rathbun, 1902 – third-stage larvae, first record in Romania.

The decapod *P. macrodactylus* has been reported since 2002 in the Romanian Black Sea, and subsequently in the Varna Lake (2009), the Kerch Strait (Sea of Azov, Chushka Spit coast, July 2018) and the Azov–Don Basin (September 2022) by several authors (Micu and Nita 2009;

Raykov et al. 2010; Timofeev et al. 2019; Matishov et al. 2022). However, no larvae had been described so far in the natural environment. Herein, we report the first finding of zoea in the third stage in two samples collected in August 2024 in the Romanian Black Sea waters. The oriental shrimp third stage larvae were distinguished by the following characters: two dorsal spines on the carapace (one clearly visible in Figure S10A), not yet developed pleopods, the sixth somite of the pleon separated from the telson and uropods, and an upward dorsal spine on telson (Figure S10B). The specimens measured about 3 mm in length.

- *Pseudodiaptomus marinus* Sato, 1913 – first record in Spanish Mediterranean waters (Algeciras Harbour).

Pseudodiaptomus marinus (Figure S11) is a calanoid copepod characteristic of coastal waters and the lower parts of estuaries (Walter 1989), native to the Northwest Pacific, from Russia to Philippines (Fleminger and Hendrix Kramer 1988). It has been widely spread throughout the Mediterranean and neighbouring waters due to ballast water and surface currents (Uttieri et al. 2023), although last westernmost record was published in the Berre Lagoon in France, a very enclosed ecosystem, in 2012 (Delpy et al. 2012). The species was later found in the Guadalquivir Estuary (Gulf of Cadiz) on the Atlantic Spanish south-western coast in 2016 (Reyes-Martínez and González-Gordillo 2019). *Pseudodiaptomus marinus* has a very successful population dynamics, caused by a continuously high egg production from May to October (Uye et al. 1982) and tolerates a wide range of temperature and salinity, from 8 °C up to 37 °C (Liang and Uye 1997) and 4 to 44 PSU (Orsi et al. 1991). The present study found *P. marinus* peaking up to 17.43 ind/m³ in two of the five sampling points of the Algeciras Commercial Harbour in April 2023 using 200 µm nets, under the Spanish Marine Strategy Framework Directive (MSFD) monitoring program (Descriptor 2: Non-indigenous species), indicating maritime transport as introduction vector. This is the first record in the Spanish Mediterranean waters and the westernmost locality recorded in the Mediterranean Sea.

- *Paracalanus tropicus* Andronov, 1977 – first record in Mediterranean waters (Algeciras Harbour, Spain).

Paracalanus tropicus (Figure S12) is an epipelagic calanoid copepod, present in coastal to offshore waters and endemic to the south Atlantic Ocean, Gulf of Guinea, Arabian Sea and Sea of Japan (Khelifi-Touhami and Ounissi 2023). *Paracalanus tropicus* is usually encompassed in the *Paracalanus parvus* Claus, 1863 species complex due to the cryptic differentiation of this group, including besides the nominal and present species also *P. indicus* Wolfenden, 1906, *P. quasimodo* Bowman, 1971, *P. nanus* Sars G.O., 1925, *P. intermedius* Shen & Bai, 1956, and *P. serrulus* Shen & Lee, 1963 (Cornils and Held 2014). Currently, taxonomic uncertainty limits species level monitoring efforts (Peters et al. 2025), usually needing genetical

analyses for ratification (Hidaka et al. 2016). Specimens of *P. tropicus* were found peaking up to 263.76 ind/m³ in three of the five sampling points of the Algeiras Commercial Harbour in September 2023 using 200 µm nets, under the Spanish MSFD monitoring program, indicating a marine transport related introduction vector. Under the microscope, the diagnostic taxonomic trait distinguishing *P. tropicus* from *P. indicus* is the prosome:urosome ratio (Pr:Ur = 3.8–4.0) (Khelifi-Touhami and Ounissi 2023). Our specimens had a mean Pr:Ur ratio of 3.86 ± 0.46 (N = 10). Together with an eDNA match for *P. tropicus* (Alvite 2023) from the adjacent harbour of Málaga, obtained during the same monitoring program, confirms the species identification. Therefore, this is the first record of this species in Mediterranean waters.

Mollusca (Gastropoda)

- *Siphonaria crenata* Blainville, 1827 – first record in Syria.

The gastropod *S. crenata* is native to the Persian Gulf (Hubendick 1944) and previous studies indicate its presence as an alien species in the eastern Mediterranean Sea (Zenetos et al. 2004), after first record in Lebanon in 1967 (Crocetta et al. 2020). The species is characterized by a solid, thick shell, decorated with 10–12 irregularly strong radial ribs alternating with irregularly thin ribs (Figure S13A). The internal furrow is pale in colour (Figure S13B) and the cone is high (Figure S13C). There is no hole in the upper part of the shell. The outer surface is creamy or rusty brown, and the inner surface is marbled white, containing brown ripples in some individuals (Figure S13A, B). Seven small individuals, 10–20 mm long (Figure S13D), were collected in August 2021 from rocky substrates in the littoral zone of Ras al-Basit, northern Syria. We acknowledge that the taxonomy of the species belonging to the family Siphonariidae is subject to considerable confusion and requires further data and analysis of its genetic characteristics (Jenkins and Köhler 2024).

- *Mitrella psilla* (Duclos, 1846) – first record in Algeria.

The gastropod *M. psilla* (Figure S14), native to West Africa, was first recorded in the Mediterranean in Tunisia (Antit et al. 2009), and then in Italian waters in the Tyrrhenian and Ionian Sea (Nappo et al. 2019; Scuderi and Villari 2022), and in Spain, indicating its continued expansion into the Basin (Martínez-Ortí et al. 2020). Its presence in the port of Algiers further demonstrates its ongoing dispersal across the Mediterranean.

Bryozoa

- *Catenicella paradoxa* Rosso, 2009 – first record in Sicily/Sicily Strait.

The bryozoan *C. paradoxa* (Figure S15A) was originally described from a few colonies collected in July 2006 on the north side of Capo Passero Island

SE Sicily. These were found encrusting two algal species typical of the Infralittoral Algae biocenosis, at a depth of 11 m beneath a cliff overhang (Rosso 2009). The species was also recently reported from three stations within the Ciclopi Islands Marine Protected Area (W Ionian Sea, E Sicily), between 5 and 26 depth, again associated with Infralittoral Algae (Rosso et al. 2019; Mytilineou et al. 2016). *Catenicella paradoxa* forms inconspicuous, glassy, and transparent colonies composed of uniserial articulated branches, and remains the only catenicellid currently known from the present-day Mediterranean (Rosso and Di Martino 2023). Rosso (2009) suggested the cryptogenic status of *C. paradoxa*, proposing four different, some potentially concurrent, hypotheses to explain its presence in the Mediterranean Sea: the possible overlooking of an inconspicuous, hidden, resident species; the local differentiation of a new taxon; the human-mediated introduction of a non-native species; or the recent arrival of an undescribed species expanding its range through the Suez Canal, potentially facilitated by warming waters and global change. The discovery of a single colony in 2014, on an artificial panel deployed at approximately 20 m depth and a few meters inside a cave in the Plemmirio Marine Protected Area near Siracusa (Ionian Sea, SE Sicily), was interpreted by Rosso et al. (2018) as potentially supporting a non-native origin. Given the species' limited and localized records in the western Ionian Sea, the finding of a single colony of *C. paradoxa* on a sponge stranded on the Marina di Ragusa beach in the winter of 2022 represents the first documented occurrence in the Sicily Strait.

- *Tricellaria inopinata* d'Hondt & Occhipinti-Ambrogi, 1985 – first record in Sicily/western Ionian Sea.

The bryozoan *T. inopinata* (Figure S15B), was described based on colonies found attached to the byssus threads of the bivalve *Mytilus galloprovincialis* Lamarck, 1819, collected from the intertidal zone of the Giudecca Canal, near central Venice (northern Adriatic Sea) in 1982 (d'Hondt and Occhipinti Ambrogi 1985; Occhipinti Ambrogi 2000). This species forms small, erect, bush-like colonies a few centimetres high, composed of jointed, slightly mineralised biserial internodes, anchored to hard substrates by rhizoids in shallow waters. Colonies grow rapidly, intertwining to form dense, whitish mats/carpets. Although described from the Mediterranean, *T. inopinata* is considered non-native species in this basin. It was likely introduced accidentally, possibly through aquaculture or ship fouling, from its presumed native range in the Pacific Ocean (Occhipinti-Ambrogi et al. 2010). Since its original description, *T. inopinata* has been recorded from several western Mediterranean localities (e.g., Ben Suissi et al. 2006; Lodola et al. 2012; Ulman et al. 2017; Ferrario et al. 2018; Fortič and Mavrič 2018; Bensari et al. 2025), with its easternmost occurrence in the Old Venetian Harbor (Heraklion, Greece) (Ulman et al. 2017). The species has also spread globally (e.g., Dyrinda et al. 2000; Johnson et al. 2012; Micael et al. 2016).

Tricellaria inopinata has been classified as invasive following monitoring of its rapid spread and increase in colony abundance and surface coverage, often to the detriment of native species that decline or disappear after its arrival and establishment. For instance, this has been documented in the Venice lagoon (Occhipinti Ambrogi 2000; Corriero et al. 2007) and in some localities along the western Atlantic coast (Johnson et al. 2012). Based on expert evaluations and direct observations, *T. inopinata* has demonstrated moderate to high impacts on biodiversity and ecosystem services, primarily through competition and fouling (Tsirintanis et al. 2022). Moreover, according to Gavira-O’Neil et al. (2016), it may act as a driver of potential invasion meltdown.

Tricellaria inopinata was found in the channel connecting the Faro Lake (located in the NE corner of Sicily) to the Ionian Sea, based on samples collected in 2006 and 2008 at about 50 cm depth. This represents the first documented record for both Sicily and the western Ionian Sea; prior to this, *T. inopinata* was only known from the eastern Ionian sector, in the Old Venetian Harbor (Heraklion, Greece) (Ulman et al. 2017). At the time of collection, the species was by far dominating the community. However, subsequent targeted surveys conducted in summer 2021, and spring 2024 and 2025, failed to detect it, suggesting that its populations may undergo regression and even local disappearance, unlike in other localities. A focused investigation is needed to determine the environmental factors responsible for this short-term success.

- *Arbopercula angulata* (Levinsen, 1909) – first record in Liguria (northernmost Tyrrhenian/western Mediterranean).

The bryozoan *A. angulata* (Figure S15C) is a non-native species in the Mediterranean. It was first reported in this basin (Rosso 1994), misidentified as *Electra tenella* (Hincks, 1880), from the Ionian coast of southeastern Sicily, where it was found on stranded plastic debris in the Gulf of Catania and on experimental plastic containers deployed in Siracusa harbour at 5 m depth in meadows dominated by *Cymodocea nodosa* (Ucria) Ascherson and *Caulerpa prolifera* (Forsskål) J.V.Lamoroux. *Arbopercula angulata* was only found on containers retrieved after deployment periods of 3 to 6 months starting from 27 January 1990. The species forms thin, circular to lobate, encrusting, usually unilaminar colonies that can extend on broad surfaces. It often shows high intracolony morphological variability. Over the past dozen years, the species, still often referred to as *E. tenella*, has been reported from additional sites, including Capo Rasocolmo near Messina (Sicily, February 2012), and the Livorno harbour (Tuscany, July 2011), both in the Tyrrhenian Sea (Thessalou - Legaki et al. 2012). More recently, it has been found on floating plastic debris collected in October 2020 off the Catalonia coast, where it was identified as *Arbopercula tenella* (Subías-Barata et al. 2022). Other records also include floating ligneous cores near the Strait of

Gibraltar, where it was reported initially as *Electra* cf. *tenella* (López de la Cuadra and García-Gómez 1994), and plastic debris stranded in July 2012 in Menorca, Balearic Islands, subsequently identified as *A. angulata* (Reverter-Gil and Souto 2025). Collectively, these records indicate that *A. angulata* may have a wide distribution in the western Mediterranean. The numerous new occurrences documented in the present study suggest a constant presence of *A. angulata* at least all-around Sicily. The observation in summer 2024 of colonies almost completely covering boat buoys in the harbour of Lerici (Liguria, Italy) represents the northernmost record of this species in both the Tyrrhenian and western Mediterranean.

Echinodermata

- *Aquilonastra burtoni* (Gray, 1840) – first record in Syria.

The starfish *A. burtoni* is widely distributed in the Indian and Pacific Oceans, from the Hawaiian Islands and Australia to Madagascar, East Africa, and the Red Sea (WoRMS Editorial Board 2025). It is a Lessepsian migrant recorded in the Levantine Sea since the 1950s at depths of 8–10 m (Tortonese 1966; Koukouras et al. 2007), that later expanded its distribution to Cyprus and the Aegean Sea (Tzomos et al. 2011). Ten specimens of this species were collected in June 2024 by divers to a depth of 2 m, from marine fouling assemblages on the concrete wall of the fishing and recreational port in the city of Tartus, southern Syria. The abundance reached 6 individuals/m², and some additional individuals were found in empty oyster shells at the same site. The specimens showed variable shapes among individuals, rather than a uniform morphology (Figure S19A). The body was asymmetrical, with a diameter of approximately 15 mm (Figure S16B). The number of arms varied between three and eight, usually seven (Figure S16A). The arms were of unequal length and finger-like, wider at the base and narrowing and rounded at the end. The colour was greenish grey on the oral side, with a large, irregular, purple-brown spot in the middle, surrounded by red spots on the basal parts of each arm (Figure S16C, D). The asymmetrical shape of all individuals in the sample supports the idea of asexual reproduction by fission (Karako et al. 2002).

Chaetognatha

- *Ferosagitta* sp. – first record in Romania.

Ferosagitta sp. (Figure S17A) is a chaetognath species found in July 2024 in three stations sampled during a survey carried out across the Romanian shelf. Both juveniles and adults have been found in three out of 22 samples collected. The species differs from *F. galerita* (Dallot, 1971), the alien species reported in the Mediterranean Sea (Terbiyik et al. 2007) by the shape of the anterior fins (narrow, elongate in our specimens) (Figure S17B), the position of the seminal vesicles (touching the posterior fins in

Ferosagitta sp.) (Figure S17C), tail shape and length (short, about 1.6 mm long, pear-like, caudal fin fan-shaped, with radiating rays, symmetric) (Figure S17C). Apart from the prominent ventral ganglion (Figure S17D), the juveniles of our species resemble the adults' general conformation. The species was found across Mangalia and Sf Gheorghe transects in near-surface and deep-water layers (0–10 m; within thermocline: 10–35 m, and below thermocline: 150 m), with densities ranging from 0.98 to 13.69 ind. m⁻³.

Tunicata (Ascidiacea)

- *Symplegma rubra* Monniot C., 1972 – first record in the Mediterranean.

Symplegma rubra (Figure S18) is a colonial ascidian with a tropical and warm-temperate distribution. In the Mediterranean, it has historically been confused with the congeners *S. brakenhielmi* (Michaelsen, 1904) and *S. viride* Herdman, 1886 (Ramos-Esplá et al. 2020). However, *S. rubra* and *S. brakenhielmi* present lobed testes in contrast to *S. viride*, while *S. brakenhielmi* presents reticulate tunic, *S. rubra* has opaque tunic and characteristic red rings encircling both siphons, although this last character has also been observed in some varieties of *S. brakenhielmi* (Herdman 1906; Tokioka 1961). Moreover, specimens identified as either *Symplegma* sp., *S. brakenhielmi*, and *S. rubra* have been found to belong to different lineages, stressing the need of taxonomic revisions based on integrated data (Mastrototaro et al. 2019). Nonetheless, the specimens, here reported from Tunisia, off the coast of the island of Kuriat, morphologically match *S. rubra* and represent the first record of this species for the Mediterranean Sea.

- *Symplegma brakenhielmi* Michaelsen, 1904 – first record in Spanish Mediterranean waters (Figure S19).

Symplegma brakenhielmi is a colonial ascidian with a pantropical distribution, present in the Mediterranean from East to West (Ramos-Esplá et al. 2020), extending in the Tyrrhenian Sea (Ulman et al. 2017; Mastrototaro et al. 2019). The colonies observed in the Port of Alicante are red and yellow (Mastrototaro et al. 2019) with a reticulated tunic, without red rings encircling both siphons, and present lobed testes.

- *Phallusia nigra* Savigny, 1816 – first record in Italy.

The solitary ascidian *P. nigra* (Figure S20), originally considered a circumtropical species, was first reported in the Mediterranean Sea by Pérès (1958), in Israel. Later, Shenkar and Loya (2009) documented a wide distribution of the species along the Israeli coasts, from shallow waters down to 30 m, on both natural and artificial substrates. The species has also been reported in Lebanon (Zibrowius and Bitar 2003), Syria (Bitar et al. 2003), along the Turkish coasts (Çinar et al. 2006), Greece (Kondilatos et al. 2010; Latsoudis 2020), and in the Aegean Sea (Thessalou-Legaki et al. 2012).

In 2024, the first record of the species in Tunisian waters was reported by Mnasri-Afifi et al. (2024a), from buoys in the port of Bizerte (northern Tunisia), representing the westernmost sighting in the Mediterranean basin. In July 2024, during hard bottom sampling (scraping) the tunicate was found in two samples collected at depths of 2 m and 5 m on the quay wall of the Levante Pier in the Port of Catania. This record represents the first morphological evidence of the black tunicate in Italian waters, whose presence in Italian seas (ports of Trieste and Civitavecchia) was previously only inferred based on eDNA metabarcoding methods (Varrella et al. 2025). Following the Catania record, the species was sampled in the port of Augusta during a Consorzio Nazionale Interuniversitario per le Scienze del Mare (CoNISMa) survey and were further confirmed by DNA barcoding by the same team (sequence deposited in GenBank under the accession code PV018849, giving a 99% identity with deposited sequences of *P. nigra*).

- *Microcosmus exasperatus* Heller, 1878 – first record in Italy.

The identity and distribution of the ascidian *M. exasperatus* (Figure S21) in the Mediterranean Sea have been the object of some confusion, as this species was historically reported as a misidentification of another, more widespread, congeneric species, *M. squamiger* Michaelsen, 1927 (Ramos-Esplá et al. 2013). Revisions of Mediterranean records of this species confirmed its occurrence in the eastern and southern Mediterranean Sea (Turon et al. 2007; Ramos-Esplá et al. 2013). The specimens sampled in the port of Augusta represent the first record for Italian waters and might have arrived either by natural expansion from Tunisia, where the species is widespread, or aided by shipping from the eastern Mediterranean Sea. A similar situation explains the occurrence of the other alien ascidian *Phallusia nigra* in the same port.

- *Distaplia bermudensis* Van Name, 1902 – first record in Italy (Adriatic Sea) and Tunisia.

Distaplia bermudensis (Figure S22) is a small colonial ascidian originally described from Bermuda and widespread in the western Atlantic. It was first reported in the Mediterranean Sea in 2000 from the Gulf of Taranto (Italy), where subsequent studies on fouling assemblages indicate an established population (Mastrototaro and Brunetti 2006). A further record in 2019 in the Phlegraean lagoons (Gulf of Naples, Tyrrhenian Sea) also pointed to an established population in that area, with the Bacoli zone as the most likely introduction site, associated with marinas and mussel farming (Virgili et al. 2022). These two Italian populations appear geographically isolated and were never compared molecularly, so they may represent separate introduction events. Both sites are associated with marinas and mussel farming, suggesting these vectors may have facilitated its introduction (Orfanidis et al. 2021; Virgili et al. 2022). In 2021,

D. bermudensis was also recorded in three Tunisian marinas (first record for Tunisia), indicating a further southward expansion of this NIS in the Mediterranean. In 2024, an extremely small colony was recorded in the Brindisi port during a marina survey, representing the first report for the Adriatic Sea. Its introduction is likely linked to shipping or aquaculture activities.

- *Botrylloides niger* Herdman, 1886 – first record in Greece (Zakynthos Island, eastern Ionian Sea).

In October 2022, the colonial ascidian *B. niger* (Figure S23), was recorded for the first time on Zakynthos Island (Ionian Sea, Greece) during a rapid assessment survey of alien species. Colonies were observed at six of nine surveyed marinas, growing on artificial substrates such as ropes and tyres, and on the basal leaves and rhizomes of the invasive seagrass *Halophila stipulacea* (Forsskål) Ascherson. The same seagrass patch also hosted dense aggregations of the bryozoan *A. verticillata*, highlighting the role of *H. stipulacea* as a novel substrate for IAS. Although *B. niger* has previously been reported as an epiphyte on the surface of native seagrasses (Virgili et al. 2022), its occurrence on the invasive *H. stipulacea* raises questions about how interactions between co-occurring IAS may shape invasion dynamics. Given the seasonal shedding of *H. stipulacea* leaves in the Mediterranean, such associations may be temporary at the colony level, but detached leaves could also facilitate short-distance dispersal by rafting. In this case, the overgrowth of *H. stipulacea* by *B. niger* could reflect a form of invasional interference, where one IAS suppresses another, potentially altering invasion trajectories and ecosystem outcomes (Simberloff and Von Holle 1999). This contrasts with invasional meltdown, in which invaders facilitate each other's spread (Simberloff and Von Holle 1999). While such interactions may limit individual IAS, cumulative impacts can still be profound due to ecosystem-level disruptions (Gordon 1998).

The ability of *B. niger* to colonize both artificial structures and living macrophytes reflects its ecological plasticity and raises concern over its potential to displace native benthic assemblages. Its presence in Zakynthos expands its known range in the Mediterranean, where it has shown high morphological and genetic variability (Virgili et al. 2022; Temiz et al. 2023; Mnasri-Afifi et al. 2024b). Alongside a rise in introduced ascidian species from 14 to 20 in the last decade in the Mediterranean Sea (Muñoz and Ramos-Esplá 2019), this finding underscores the need for early detection, molecular surveillance, and ecosystem-based management in addressing marine bioinvasions. For example, the presence of *B. niger* has been previously hypothesized in Crete, South Aegean Sea, as early as 2018, on Artificial Reef Monitoring Structures (ARMS) via molecular screening, as manifested in the accompanying dataset of Obst et al. (2020). On the other hand, the analysis of *B. niger* colonies from the Mediterranean Sea has

already proved to be a potential biological resource for nutritional or functional purposes, reducing, in this way, the economic losses caused by this invasive organism (Della Sala et al. 2022). Therefore, its potential use for such purposes can be also tested in the new localities of occurrence reported here.

Macroalgae (Ochrophyta)

- *Stypopodium schimperi* (Kützinger) Verlaque & Boudouresque – First record in southern Italy (Lampedusa & Pantelleria).

Stypopodium schimperi is a medium sized brown macroalga (up to 18 cm high) irregularly branched with fan-shaped blades (Figure S24). Its distinguishing characteristics are the flat growing margin, the absence of calcification, the hairs in concentric lines and the distinctive tetrasporangia external to the blade (Verlaque et al. 2015). It is distributed in shallow subtidal coral and rocky reefs in the Indian and Atlantic Oceans. It has been introduced in the Mediterranean Sea, most probably through the Suez Canal. It was first recorded in Israel in 1973 (Lundberg 1980) and is now very common in the Eastern Mediterranean with records in Libya (Nizamuddin and Godeh 1989), Syria (Mayhoub 1989), Egypt, Turkey, Cyprus (Verlaque and Boudouresque 1991), Lebanon (Bitar et al. 2000), Greece (Sartoni and De Biasi 1999; Tsiamis et al. 2008; Nicolaidou et al. 2012) and recently Croatia and Adriatic Sea (Žuljević et al. 2023). In summer 2024, it was observed for the first time in Southern Italy, in the islands of Lampedusa and Pantelleria. It is likely that the species is more widely distributed in the Mediterranean and maybe confused with *Lobophora variegata* (J.V. Lamouroux) Womersley ex E.C. Oliveira another brown algae with a similar morphology.

Discussion

This third collaborative compilation of unpublished records further consolidates an open, region-wide evidence base for alien, cryptogenic, and neonative species in the Mediterranean and Black Sea. By adding 7,434 geo-referenced records from 21 countries, it extends the temporal span (1973–2025) and complements two earlier datasets (Katsanevakis et al. 2020; Ragkousis et al. 2023) that collectively demonstrated the utility of mobilizing otherwise inaccessible information held by experts and institutions. Together, these efforts show that routine aggregation of unpublished records can rapidly enrich inventories, improve the spatial and taxonomic resolution of knowledge, and enable timely reassessments of invasion dynamics needed for policy and management (Katsanevakis et al. 2020; Ragkousis et al. 2023).

The spatiotemporal distribution of records highlights both strengths and gaps in current monitoring. Contributions were highest in Lebanon, Greece, Cyprus, Italy, and Romania, while Albania, Egypt, Georgia, Ukraine, France,

and parts of North Africa remained underrepresented. Similar geographic imbalances have been reported in the previous two “unpublished records” datasets, suggesting that uneven expert participation/sampling effort (alongside ecology) systematically shapes apparent hotspots and gaps (Katsanevakis et al. 2020; Ragkousis et al. 2023). This underrepresentation should not be interpreted exclusively as a lack of national expertise or monitoring programme; for example, Massé et al. (2023) provide an updated overview of marine alien (including cryptogenic) species across three metropolitan French regions. At the basin scale, expert-validated inventories (e.g., Galanidi et al. 2023) provide policy-oriented baselines, which our record-level, georeferenced compilation can complement by adding finer spatiotemporal resolution that supports inventory updates and monitoring prioritization.

Habitat was reported for a subset of records; within those, 453 records explicitly indicate observations in harbours/marinas or shellfish aquaculture settings (e.g., mussel/oyster farming). These represent a minimum count and are consistent with the role of artificial coastal infrastructures and aquaculture facilities as introduction gateways and detection hotspots for alien species. Because habitat/site type was not reported for all entries and does not allow a standardized classification of all records by site category, we do not attempt a full quantitative partition of the dataset by site type.

Temporally, reporting increased sharply after 2021, peaking in 2024 – a trend shaped not only by species spread but also by intensified surveys and improved contributor engagement. Such patterns caution against interpreting temporal increases as purely ecological signals. Instead, they flag locations where targeted effort, training, and data-sharing could most improve coverage (see also Katsanevakis et al. 2020; Ragkousis et al. 2023).

Taxonomic uncertainty is inevitable in a large, multi-source dataset spanning many groups and a large number of contributors, and some misidentifications are likely – particularly for taxa that require specialist examination or integrative taxonomy (e.g., cryptic species complexes and closely related congeners). Records should therefore be interpreted as best-available expert identifications at the time of reporting, and they may be refined in future as taxonomic knowledge improves, and as additional voucher or molecular evidence becomes available (see also Zenetos et al. 2017). For transparency and traceability, the identifier is reported for every record in Table S1, responsibility for correct identification and reporting rests with the observer of each record, as in previous iterations of this collective data initiative.

Several new basin-level records (e.g., *D. amblysinus*, *S. rubra*, *M. mcintoshii*, *P. tropicus*) and national firsts (e.g., in Algeria, Cyprus, Greece, Italy, Slovenia, Spain, Syria, Tunisia) show that introductions continue across taxa and pathways. Range extensions (e.g., *P. depauperata* in the Tyrrhenian; *A. angulata* northward; *S. brakenhielmi* westward) illustrate secondary

spread, often linked to shipping, marinas, or aquaculture. The isolated Italian populations of *D. bermudensis* highlight how repeated introductions may mimic continuity if molecular comparisons are lacking, while *P. marinus* in Algeciras emphasizes ballast water as a recurrent vector.

A complementary dimension not analysed quantitatively here is the donor-region structure of introductions (e.g., Atlantic vs Red Sea/Indo-Pacific sources), a topic discussed at basin scale by Tsiamis et al. (2018). Robust estimates would require standardized species-level attribution of native range for primary and secondary introductions at subregional level, which is not consistently captured as a structured field in Table S1 and would involve additional harmonization and literature-based validation for many taxa. We therefore treat donor-region partitioning as a priority for a subsequent synthesis, ideally integrating the three unpublished-record compilations with published inventories and origin/pathway classifications.

The dataset also sheds light on ecological dynamics. Cryptogenic cases such as *M. mcintoshii* illustrate how symbiotic lifestyles complicate origin assessments. Neonatives are expected to continue to expand under climate change, while interactions among invaders (e.g., *B. niger* overgrowing *H. stipulacea*) raise questions about “invasional interference” versus facilitation. These examples underline the need for an ecosystem-based perspective in monitoring and management, especially in port and marina environments.

As in earlier compilations, biases in sampling effort and taxonomic expertise limit quantitative inference, but do not reduce the dataset’s value for early detection, range tracking, and prioritization of surveillance, including targeted molecular confirmation where appropriate. Continued coordination, standardization, and integration with official databases remain essential. Overall, this initiative confirms that collaborative mobilization of unpublished records is a cost-effective way to sharpen regional situational awareness and guide proactive responses to marine bioinvasions.

This work demonstrates how sustained collaboration and systematic mobilization of unpublished records can fill critical knowledge gaps on alien, cryptogenic, and neonative species in the Mediterranean and Black Seas. By improving the spatiotemporal resolution of regional inventories, the dataset supports early detection and prioritization of monitoring, contributing to EU-level obligations under the Marine Strategy Framework Directive (MSFD 2008/56/EC)—including Descriptor 2 (non-indigenous species)—and facilitating interoperability with MSFD-linked infrastructures such as EASIN (European Alien Species Information Network). It also supports region-wide commitments under the Barcelona Convention and UNEP/MAP, including the updated Action Plan concerning Species Introductions and Invasive Species in the Mediterranean Sea adopted in 2023 (Decision IG.26/5, Annex IV), which emphasizes coordinated monitoring and data sharing. In addition, regional practitioner networks such as

MedPAN can provide an operational pathway to strengthen ongoing acquisition and updating of alien-species occurrence information across Mediterranean MPAs.

Continued iterations of this effort will be essential to keep pace with ongoing introductions and shifting species distributions in a rapidly changing sea.

Authors' contribution

JCAP, SK research conceptualization; data analysis and interpretation; original draft; writing – review and editing; SK: funding provision; all authors: investigation and data collection, writing – review and editing.

Author ORCIDs (where available)

Jennifer C.A. Pistevos: [0000-0001-8081-7069](https://orcid.org/0000-0001-8081-7069); Argyro Zenetos: [0000-0003-2670-042X](https://orcid.org/0000-0003-2670-042X); Yolanda Stranga: [0009-0003-4231-856X](https://orcid.org/0009-0003-4231-856X); Athanasios Nikolaou: [0000-0003-2241-3984](https://orcid.org/0000-0003-2241-3984); Joachim Langeneck: [0000-0003-3665-8683](https://orcid.org/0000-0003-3665-8683); Mihaela Muresan: [0000-0002-8446-1263](https://orcid.org/0000-0002-8446-1263); Juan Sempere-Valverde: [0000-0001-5856-9214](https://orcid.org/0000-0001-5856-9214); Sahar Chebaane: [0000-0003-3700-8930](https://orcid.org/0000-0003-3700-8930); Lydia Png-Gonzalez: [0000-0001-7986-9654](https://orcid.org/0000-0001-7986-9654); Sylvaine Giakoumi: [0000-0002-3301-5538](https://orcid.org/0000-0002-3301-5538); Michail Ragkousis: [0000-0002-3886-4280](https://orcid.org/0000-0002-3886-4280); Adriana Vella: [0000-0003-2246-5354](https://orcid.org/0000-0003-2246-5354); Noel Vella: [0000-0003-0080-5709](https://orcid.org/0000-0003-0080-5709); Vesselina Vasileva Mihneva: [0000-0003-4581-7738](https://orcid.org/0000-0003-4581-7738); Thanos Dailianis: [0000-0002-4102-9331](https://orcid.org/0000-0002-4102-9331); Victor Surugiu: [0000-0003-4933-7678](https://orcid.org/0000-0003-4933-7678); Angeliki Bosmali: [0009-0003-4767-0300](https://orcid.org/0009-0003-4767-0300); Vasilis Gerovasileiou: [0000-0002-9143-7480](https://orcid.org/0000-0002-9143-7480); Fabio Crocetta: [0000-0002-2800-3735](https://orcid.org/0000-0002-2800-3735); Leda Liyue Cai: [0000-0001-7699-4601](https://orcid.org/0000-0001-7699-4601); Angela G. Bartolo: [0000-0001-9109-8979](https://orcid.org/0000-0001-9109-8979); Maria Rousou: [0000-0003-1234-2027](https://orcid.org/0000-0003-1234-2027); Jose Antonio García-Charton: [0000-0002-5973-9913](https://orcid.org/0000-0002-5973-9913); Sandra Agius Darmanin: [0000-0003-4489-9030](https://orcid.org/0000-0003-4489-9030); Riccardo Scalera: [0000-0002-4100-3959](https://orcid.org/0000-0002-4100-3959); Donat Petricioli: [0000-0002-0150-9322](https://orcid.org/0000-0002-0150-9322); Tatjana Bakran-Petricioli: [0000-0002-3368-4248](https://orcid.org/0000-0002-3368-4248); Alberto Colletti: [0000-0002-7027-6315](https://orcid.org/0000-0002-7027-6315); Stefanos Kalogiour: [0000-0002-3064-9236](https://orcid.org/0000-0002-3064-9236); Domenico Sacco: [0009-0003-6595-1931](https://orcid.org/0009-0003-6595-1931); Tullio Rossi: [0000-0003-0461-554X](https://orcid.org/0000-0003-0461-554X); Christophe Lejeusne: [0000-0002-8821-3849](https://orcid.org/0000-0002-8821-3849); Dominique Davoult: [0000-0001-8678-3998](https://orcid.org/0000-0001-8678-3998); Daniel Faget: [0000-0001-7122-7446](https://orcid.org/0000-0001-7122-7446); Petros Stavrou: [0009-0006-0670-0237](https://orcid.org/0009-0006-0670-0237); Jasmine Ferrario: [0000-0003-4462-4141](https://orcid.org/0000-0003-4462-4141); Stelios Katsanevakis: [0000-0002-5137-7540](https://orcid.org/0000-0002-5137-7540); Arthur R. Bos: [0000-0002-0708-314X](https://orcid.org/0000-0002-0708-314X); Grigorios Skouradakis: [0000-0002-8482-460X](https://orcid.org/0000-0002-8482-460X); Konstantinos Kavakakis: [0009-0003-0583-8304](https://orcid.org/0009-0003-0583-8304); Carlo Nike Bianchi: [0000-0003-1071-3285](https://orcid.org/0000-0003-1071-3285); Carla Morri: [0000-0003-4520-2741](https://orcid.org/0000-0003-4520-2741); Laura Tamburello: [0000-0003-1305-8802](https://orcid.org/0000-0003-1305-8802); Bessy Stancanelli: [0000-0001-7126-0633](https://orcid.org/0000-0001-7126-0633); Vincenzo Di Martino: [0000-0002-5987-0954](https://orcid.org/0000-0002-5987-0954); Julian Evans: [0000-0001-7837-5927](https://orcid.org/0000-0001-7837-5927); Abigail Kennedy: [0009-0006-8924-7588](https://orcid.org/0009-0006-8924-7588); Daniele Grech: [0000-0002-4936-5997](https://orcid.org/0000-0002-4936-5997); Maria Zotou: [0000-0002-9495-3907](https://orcid.org/0000-0002-9495-3907); Ernesto Azzurro: [0000-0002-9805-3164](https://orcid.org/0000-0002-9805-3164); Patrick J Schembri: [0000-0002-6723-7198](https://orcid.org/0000-0002-6723-7198); Ilaria D'Aniello: [0009-0001-7720-2335](https://orcid.org/0009-0001-7720-2335); Michele Solca: [0000-0002-3190-5844](https://orcid.org/0000-0002-3190-5844); Antonios Mazaris: [0000-0002-4961-5490](https://orcid.org/0000-0002-4961-5490); Frédérique Viard: [0000-0001-5603-9527](https://orcid.org/0000-0001-5603-9527); Michele Luca Geraci: [0000-0002-3143-4659](https://orcid.org/0000-0002-3143-4659); Giulia Furfaro: [0000-0001-8184-2266](https://orcid.org/0000-0001-8184-2266); Giacomo Sardo: [0000-0001-7997-637X](https://orcid.org/0000-0001-7997-637X); Sergio Vitale: [0000-0001-6063-4126](https://orcid.org/0000-0001-6063-4126); Fabio Falsone: [0000-0003-1754-4226](https://orcid.org/0000-0003-1754-4226); Monica Calabrò: [0000-0002-9972-0440](https://orcid.org/0000-0002-9972-0440); Louis Hadjioannou: [0000-0001-5864-5467](https://orcid.org/0000-0001-5864-5467); Michel Bariche: [0000-0001-6831-4311](https://orcid.org/0000-0001-6831-4311); Alfonso A. Ramos Esplá: [0000-0002-6886-218X](https://orcid.org/0000-0002-6886-218X); Luca Castriota: [0000-0002-3349-9501](https://orcid.org/0000-0002-3349-9501); Manuela Falautano: [0000-0001-7844-665X](https://orcid.org/0000-0001-7844-665X); Ana Fortič: [0000-0002-5715-3572](https://orcid.org/0000-0002-5715-3572); Borut Mavrič: [0000-0001-5488-0793](https://orcid.org/0000-0001-5488-0793); Domen Trkov: [0000-0002-6573-2822](https://orcid.org/0000-0002-6573-2822); Lovrenc Lipej: [0000-0002-7608-1631](https://orcid.org/0000-0002-7608-1631); Martina Orlando-Bonaca: [0000-0001-7700-7663](https://orcid.org/0000-0001-7700-7663); Valentina Pitacco: [0000-0003-2724-2838](https://orcid.org/0000-0003-2724-2838); Neža Leban: [0009-0009-5913-2693](https://orcid.org/0009-0009-5913-2693); Antonietta Rosso: [0000-0001-5565-9513](https://orcid.org/0000-0001-5565-9513); Emanuela Di Martino: [0000-0002-3892-4036](https://orcid.org/0000-0002-3892-4036); Gemma Donato: [0000-0003-1783-4688](https://orcid.org/0000-0003-1783-4688); Rossana Sanfilippo: [0000-0001-6154-8027](https://orcid.org/0000-0001-6154-8027); Chiara Siddiolo: [0000-0003-1871-5955](https://orcid.org/0000-0003-1871-5955); Rym Zakhama-Sraieb: [0000-0002-5174-9789](https://orcid.org/0000-0002-5174-9789); Intissar Mnasri: [0009-0004-7609-7950](https://orcid.org/0009-0004-7609-7950); Víctor Orenes-Salazar: [0000-0001-9291-1087](https://orcid.org/0000-0001-9291-1087); Antonio Di Franco: [0000-0003-3411-7015](https://orcid.org/0000-0003-3411-7015); Anna Maria Mannino [0000-0001-6419-7066](https://orcid.org/0000-0001-6419-7066).

Acknowledgements

On 14 May 2026, whilst this article was still in press, our co-author, Professor Monica Montefalcone, passed away at the age of 52 following a diving accident. Monica was a highly respected Italian researcher and a wonderful person, whose loss is deeply felt by the entire marine science community. The authors dedicate this work to her memory.

Professor Ghazi Bitar, in addition to his personal records, achieved many others during Lebanese-French cooperation missions (CEDRE Project between the Lebanese University, the Lebanese CNRS, and IMBE UMR CNRS7263 / IRD 237 / Aix Marseille Université, Station

Marine d'Endoume), Lebanese-Italian cooperation (with Fabio Crocetta, Marco Oliverio and Francesco Tiralongo), and also within the framework of several national projects in close collaboration with the SPA/RAC in Tunis (MedMPAnet project), as well as with the University of Alicante and OCEANA.

The data from ARPA SICILIA (Paolo Balistreri, Eleonora Macaluso, and Benedetto Sirchia) were produced thanks to the Operational Agreement MASE-ISPRA-ARPA of February 16, 2024, implementing Legislative Decree No. 190/2010 transposing Directive 2008/56/EC (Marine Strategy Framework Directive) for the 2024–2026 period. Monitoring Activity Module 3, Marine Strategy Framework Directive 2008/56/EC.

The biologists wish to thank the following for their collaboration in identifying the species *Phallusia nigra* Savigny, 1816: Dr. Intissar Mnasri-Afifi (University of Tunis El Manar) and Dr. Alfonso A. Ramos Esplá (University of Alicante); and for the identification of *Dynoides amblysinus* Pillai, 1954: Drs. Niel Bruce (Queensland Museum), Panakkool Thamban Aneesh (Hiroshima University), Ghazi Bitar (Lebanese University), and José Castelló (University of Barcelona).

Vasilis Gerovasileiou, Thanos Dailianis and Markos Digenis received support for sampling activities by (i) the Hellenic Foundation for Research and Innovation (HFRI) under the 4th Call for HFRI PhD Fellowships (Fellowship Number: 10597); (ii) the project “Study and clean-up activities of marine caves and rapid assessment of pressures/threats, non-indigenous species and protected benthic species”, in the framework of the action “Support of the Dodecanese Protected Areas Management Body for the implementation of management measures for protected areas, species and habitats (MIS 5034797)”, of the NSRF 2014–2020 Operational Program “Transport Infrastructure, Environment and Sustainable Development”; and (iii) the project “SponBIODIV - Marine sponge biodiversity from genes to ecosystems: delivering knowledge and tools for sustainable management and conservation”, funded by Biodiversa+, the European Biodiversity Partnership under the 2021–2022 BiodivProtect joint call for research proposals, co-funded by the European Commission (GA N°101052342) and the funding organisation General Secretariat for Research and Innovation (GSRI) under the Hellenic Ministry of Development (ITBIO-0559429). They also thank the Management Units of the Southeastern Aegean Protected Areas, and Zakynthos and Ainos National Parks and Protected Areas of the Ionian islands of the Natural Environment and Climate Change Agency (NECCA) for providing sampling permissions; Dennis Mohr and the staff of Nero Sport Diving Center (Limni Keriou, Zakynthos) and Dinos Protopapas from the Karpathos Diving Center (Pigadia, Karpathos) for their valuable knowledge and help during sampling in Zakynthos, Karpathos and Saria Islands, respectively.

Antonietta Rosso, Emanuela Di Martino, Gemma Donato and Rossana Sanfilippo received funds from the University of Catania, through PiaCeRi—PIAno inCEntivi per la Ricerca di Ateneo—2024 – 26 – Linea di intervento 1 – Ricerca Collaborativa – Progetto COMBIO, and this paper is the Catania Palaeontological Research Group contribution n. 537.

T.E. Kampouris and Z. Erga wish to warmly thank the following individuals: Nikos Lazaridis and Lazaros Lazaridis, professional coastal fishermen for providing information on the *P. aztecus* population status at Thermaikos Gulf, NW Aegean Sea; from Dodecanese: Fokas Evgenikos, spearfisher, Stavros Melianos, professional fisherman, and Grigoris Ergas amateur fisherman for informing us on the occurrence of several *S. ghobban* specimens.

The work of the researchers from the Marine Biology Station Piran (National Institute of Biology) was possible thanks to the financial support of the Ministry of Natural Resources and Spatial Planning of the Republic of Slovenia within implementation of WFD and MSFD, of the Agency for Environment within national monitoring of Ecological status of coastal waters within the Republic of Slovenia, and of the Slovenian Research Agency (research core funding No. P1-0237).

A. Carbonell, JM Cañizares-González, L. Png-Gonzalez, R. Comas-González and M. Vázquez-Luis were funded by ESMARES3 project “Monitoring Marine Strategies, Assessment of the Marine Environment, and Definition of Good Environmental Status”, carried out by the IEO-CSIC as part of the Interdepartmental Collaboration Agreement between the Ministry for Ecological Transition and Demographic Challenge, and the Ministry of Science, Innovation, and Universities; co-funded by the European Maritime, Fisheries, and Aquaculture Fund.

M. Salomidi, V. Gerakaris, Y. Issaris, E. Kytinou, P. Lardi, G. Oikonomidis and K. Tsirintanis collected their provided records during fieldwork of the Hellenic Centre for Marine Research (HCMR) for the implementation of the European Directives HD 92/43/EC, WFD (2000/60/EC) and MSFD (2008/56/EC).

Prof. Dr. Ergün Taşkin was supported by TÜBİTAK (project number: 121Y215).

C.G. Georgiadis samples were identified during sampling by the Laboratory of Benthic Ecology of the Fisheries Research Institute at the port of Volos, as part of the work of the Marine Strategy Framework Directive, led by Dr. Sotiris Orfanidis.

This publication was produced while Chiara Siddiolo is attending the PhD programme in Sustainable Development And Climate Change at the University School for Advanced Studies IUSS Pavia, Cycle XXXVIII, with the support of a scholarship financed by the Ministerial

Decree no. 351 of 9th April 2022, based on the NRRP— funded by the European Union— NextGenerationEU—Mission 4 “Education and Research”, Component 1 “Enhancement of the offer of educational services: from nurseries to universities” -Investment 4.1 “Extension of the number of research doctorates and innovative doctorates for public administration and cultural heritage”, funded by the University of Catania.

Prof. Monica Montefalcone, Dr. Annalisa Azzola and Prof. Anna Maria Mannino activities were partially funded under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 - Call for tender No. 3138 of 16 December 2021, rectified by Decree n.3175 of 18 December 2021 of Italian Ministry of University and Research funded by the European Union – NextGenerationEU; Award Number: Project code CN_00000033, Concession Decree No. 1034 of 17 June 2022 adopted by the Italian Ministry of University and Research, CUP D33C22000960007, Project title “National Biodiversity Future Center - NBFC”.

María Altamirano collected her records funded by the projects RUGULOPTERYX (Fundación Biodiversidad 2020) and FEDERJA-006 (FEDER-Junta de Andalucía).

Dr. Christophe Lejeune and Dr. Daniel Faget received support from the French government under the France 2030 investment plan, as part of the Initiative d’Excellence d’Aix-Marseille Université - A*MIDEX (AMX-21-RID-036).

Daniele Grech was supported by the “ALIEM VIGIL - Actions de limitation, de gestion et de vigilance liées à l’introduction d’espèces exotiques envahissantes dans les zones de transports en Méditerranée”, project financed under the Interreg Italia - France Maritime 2021–2027 (<https://www.aliem-network.eu/>).

Dr. Sahar Chebaane was financially supported by doctoral fellowships by Agência Regional para o Desenvolvimento da Investigação, Tecnologia e Inovação (ARDITI-M1420-09-5369-FSE-000002). Dr. Juan Sempere-Valverde was supported by a FPI Grant (PRE2018-086266) from Ministerio de Ciencia, Innovación y Universidades (Project CGL 2017-82739-P) co-financed by ERDF European Union and Agencia Estatal de Investigación, Gobierno de España.

Prof. Carlos Navarro-Barranco benefited from the mobility grant Ayudas de Movilidad para Estancias en el Extranjero. Plan Propio. University of Seville. We would like to thank the Regional Activity Center for Specially Protected Areas (RAC/SPA) for supporting JSV and SC surveys in Tunisia.

Carlos Navarro-Barranco was financially supported by the grant “Ayudas de Movilidad para Estancias en el Extranjero. Plan Propio,” awarded by the University of Seville.

The research was supported by the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 - Call for tender No. 3138 of 16 December 2021, rectified by Decree n.3175 of 18 December 2021 of Italian Ministry of University and Research funded by the European Union— NextGenerationEU; Award Number: Project code CN_00000033, Concession Decree No. 1034 of 17 June 2022 adopted by the Italian Ministry of University and Research, CUP C63C22000520001, Project title “National Biodiversity Future Center - NBFC”; PRIN 2022 project Reconnect “Reconciling conservation and exploitation of a keystone species through networks of Marine Protected Areas” (CUP C53D23003550006), funded by European Union-Next Generation EU; FishMPABlue2 and FishMPABlue2 Plus projects (<https://fishmpablue-2.interreg-med.eu/>) funded by the European Territorial Cooperation Programme MED and co-financed by the European Regional Development Fund (ERDF); “Analysis of the state of fish stocks and assessment of biomass in Telašćica nature park”, agreement no: 66/2023/hr00085 and “Analysis of the state of fish stocks and assessment of biomass in the proposed special reserve St. Stupišće – vis island” by method of visual census agreement no: 39/2024/hr00092 between Stazione Zoologica and WWF Adria.

Funding declaration

This article was supported by the European Union’s Horizon Europe HORIZON-CL6-2024-BIODIV-01 project “GuardIAS - Guarding European from IAS”, under grant agreement no. 101181413 (Katsanevakis et al. 2024). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Data availability

Species georeferenced records are available at the European Alien Species Information Network: <https://easin.jrc.ec.europa.eu/easin/RJD/Download/ca406ad8-fa04-42cf-825d-03232be3570d>

References

- Aguado MT, San Martin G (2007) Syllidae (Polychaeta) from Lebanon with two new reports for the Mediterranean Sea. *Cahiers de Biologie Marine* 48: 207–224
- Albano PG, Steger J, Bošnjak M, Dunne B, Guifarro Z, Turapova E, Hua Q, Kaufman DS, Rilov G, Zuschin M (2021) Native biodiversity collapse in the eastern Mediterranean. *Proceedings of the Royal Society B* 288: 20202469, <https://doi.org/10.1098/rspb.2020.2469>

- Antit M, Gofas S, Azzouna A (2009) A gastropod from the tropical Atlantic becomes an established alien in the Mediterranean. *Biological Invasions* 12: 991–994, <https://doi.org/10.1007/s10530-009-9532-2>
- Azzurro E, Smeraldo S, Minelli A, D'Amen M (2022) ORMEF: a Mediterranean database of exotic fish records. *Scientific Data* 9: 363, <https://doi.org/10.1038/s41597-022-01487-z>
- Ba-Akdah MA, Satheesh S, Al-Sofyani AMA, Lucas Y, Álvarez-Campos P, San Martín G (2018) Taxonomy of some species of the genus *Syllis* (Annelida: Syllidae: Syllinae) from the Red Sea found among the first colonizers of an artificial substrate. *Marine Biology Research* 14: 790–805, <https://doi.org/10.1080/17451000.2018.1531133>
- Băncilă RI, Skolka M, Ivanova P, Surugiu V, Stefanova K, Todorova V, Zenetos A (2022) Alien species of the Romanian and Bulgarian Black Sea coast: state of knowledge, uncertainties, and needs for future research. *Aquatic Invasions* 17: 353–373, <https://doi.org/10.3391/ai.2022.17.3.02>
- Barnich R, Dietrich A, Hager T, Fiege D (2017) On the genera *Malmgrenia* McIntosh, 1874 and *Pettibonesia* Nemésio, 2006 in the Northeast Atlantic and Mediterranean Sea, with descriptions of two new species (Polychaeta: Polynoidae). *Marine Biodiversity* 49: 315–324, <https://doi.org/10.1007/s12526-017-0802-4>
- Bastida-Zavala JR, ten Hove HA (2002) Revision of *Hydroides* Gunnerus, 1768 (Polychaeta: Serpulidae) from the Western Atlantic region. *Beaufortia* 52: 103–178
- Berkeley E, Berkeley C (1941) On a collection of Polychaeta from Southern California. *Bulletin of the Southern California Academy of Sciences* 40(1): 16–60
- Ben Souissi J, Ben Salem M, Zaouali J (2006) *Tricellaria inopinata* (Bryozoa, Cheilostome), récolté pour la première fois en Tunisie (Lagune sud de Tunis: Tunisie septentrionale). In: Actes de la 8ème Conférence Internationale des Limnologues d'Expression Française, Hammamet (Tunisie), 17–21 Mars 2006, pp 17–21
- Bensari B, Bahbah L, Kacimi A, Bachetarzi R, Bachari NEI (2025) Non-indigenous and cryptogenic marine species in the Port of Algiers (Western Mediterranean). *Mediterranean Marine Science* 26: 592–603, <https://doi.org/10.12681/mms.39689>
- Bianchi CN, Morri C (2003) Global sea warming and ‘tropicalization’ of the Mediterranean Sea: biogeographic and ecological aspects. *Biogeographia* 24: 319–327, <https://doi.org/10.21426/B6110129>
- Bitar G, Harmelin JG, Verlaque M, Zibrowius H (2000) Sur la flore marine benthique supposée Lessepsienne de la côte libanaise. Cas particulier de *Stypopodium schimperii*. In: Symposium On Marine Vegetation (Ajaccio, 3–4 October 2000), p 97, https://www.rac-spa.org/sites/default/files/doc_vegetation/actes_1er_sympos_veg_2000.pdf
- Boltacheva NA, Lisitskaya EV, Lebedovskaya MV (2011) New species *Hydroides dianthus* (Verrill, 1873) (Polychaeta: Serpulidae) from the coastal waters of Crimea (the Black Sea). *Morskoi ekologichnyi Zhurnal* 10(2): 34–38
- Bruce NL (1982) The genus *Dynoides* Barnard, (1914) (Crustacea: Isopoda: Sphaeromatidea) from eastern Australia, with description of new species. *Memoirs of the Queensland Museum* 20: 443–452, <https://biostor.org/reference/206678>
- Bruce NL, Wong HPS (2015) An overview of the marine Isopoda (Crustacea) of Singapore. *Raffles Bulletin of Zoology Supplement* 31: 152–168
- Carlton JT (1996) Biological invasions and cryptogenic species. *Ecology* 77: 1653–1655, <https://doi.org/10.2307/2265767>
- Çinar ME (2005) *Syllis ergeni*: a new species of Syllidae (Annelida: Polychaeta) from Izmir Bay (Aegean Sea, eastern Mediterranean Sea). *Zootaxa* 1036: 43–54, <https://doi.org/10.11646/zootaxa.1036.1.4>
- Çinar ME (2009) Alien polychaete species (Annelida: Polychaeta) on the southern coast of Turkey (Levantine Sea, eastern Mediterranean), with 13 new records for the Mediterranean Sea. *Journal of Natural History* 53: 2283–2328, <https://doi.org/10.1080/00222930903094654>
- Çinar ME (2013) Alien polychaete species worldwide: current status and their impacts. *Journal of the Marine Biological Association of the United Kingdom* 93: 1257–1278, <https://doi.org/10.1017/S0025315412001646>
- Çinar ME, Ergen Z (2007) The presence of *Chaetozona corona* (Polychaeta: Cirratulidae) in the Mediterranean Sea: an alien or a native species. *Cahiers de Biologie Marine* 48(4): 339–346
- Çinar ME, Bilecenoglu M, Öztürk B, Can A (2006) New record of alien species on the Levantine coast of Turkey. *Aquatic Invasions* 1: 84–90, <https://doi.org/10.3391/ai.2006.1.2.6>
- Çinar ME, Katağan T, Koçak F, Öztürk B, Ergen Z, Kocatas A, Önen M, Kırkim F, Bakir K, Kurt G, Dağlı E, Açık S, Doğan A, Özcan T (2008) Faunal assemblages of the mussel *Mytilus galloprovincialis* in and around Alsancak Harbour (Izmir Bay, eastern Mediterranean) with special emphasis on alien species. *Journal of Marine Systems* 71: 1–17, <https://doi.org/10.1016/j.jmarsys.2007.05.004>
- Çinar ME, Bakir K, Öztürk B, Katağan T, Doğan A, Açık S, Kurt-Şahin G, Özcan T, Dağlı E, Bitlis-Bakir B (2017) Macrobenthic fauna associated with the invasive alien species *Brachidontes pharaonis* (Mollusca: Bivalvia) in the Levantine Sea (Turkey). *Journal of the Marine Biological Association of the United Kingdom* 97: 613–628, <https://doi.org/10.1017/S0025315417000133>
- Çinar ME, Bakir K, Öztürk B, Doğan A, Açık S, Kırkim F, Dağlı E, Kurt G, Evcen A, Koçak F, Bitlis B (2020) Spatial distribution pattern of macroinvertebrates associated with the black mussel *Mytilus galloprovincialis* (Mollusca: Bivalvia) in the Sea of Marmara. *Journal of Marine Systems* 211: 103402, <https://doi.org/10.1016/j.jmarsys.2020.103402>

- Çinar ME, Bilecenoglu M, Yokeş MB, Öztürk B, Taşkın E, Bakır K, Doğan A, Açık S (2021) Current status (as of end of 2020) of marine alien species in Turkey. *PLoS ONE* 16: e0251086, <https://doi.org/10.1371/journal.pone.0251086>
- Çinar ME, Açık S, Kurt G, Erdoğan-Dereli D (2024) Diversity of Annelida from the coasts of Türkiye. *Turkish Journal of Zoology* 48: 413–335, <https://doi.org/10.55730/1300-0179.3193>
- Cornils A, Held C (2014) Evidence of cryptic and pseudocryptic speciation in the *Paracalanus parvus* species complex (Crustacea, Copepoda, Calanoida). *Frontiers in Zoology* 11: 19, <https://doi.org/10.1186/1742-9994-11-19>
- Corriero G, Longo C, Mercurio M, Marchini A, Occhipinti-Ambrogi A (2007) Porifera and Bryozoa on artificial hard bottoms in the Venice Lagoon: Spatial distribution and temporal changes in the northern basin. *Italian Journal of Zoology* 74: 1, <https://doi.org/10.1080/11250000601084100>
- Crocetta F, Bitar G, Zibrowius H, Oliverio M (2020) Increase in knowledge of the marine gastropod fauna of Lebanon since the 19th century. *Bulletin of Marine Science* 96: 1–22, <https://doi.org/10.5343/bms.2019.0012>
- d'Hondt JL, Occhipinti Ambrogi A (1985) *Tricellaria inopinata*, n. sp., un nouveau bryozoaire Cheilostome de la faune méditerranéenne. *Marine Ecology* 6: 35–46, <https://doi.org/10.1111/j.1439-0485.1985.tb00319.x>
- Dağlı E, Çinar ME (2009) Species of the subgenera *Aquilaspio* and *Prionospio* (Polychaeta: Spionidae: *Prionospio*) from the southern coast of Turkey (Levantine Sea, eastern Mediterranean), with description of a new species and two new reports for the Mediterranean fauna. *Zootaxa* 2275: 1–20, <https://doi.org/10.11646/zootaxa.2275.1.1>
- Della Sala G, Coppola D, Virgili R, Vitale GA, Tanduo V, Teta R, Crocetta F, de Pascale D (2022) Untargeted metabolomics yields insights into the lipidome of *Botrylloides niger* Herdman, 1886, an ascidian invading the Mediterranean Sea. *Frontiers in Marine Science* 9: 865751, <https://doi.org/10.3389/fmars.2022.865751>
- Delpy F, Pagano M, Blanchot J, Carlotti F, Thibault-Botha D (2012) Man-induced hydrological changes, metazooplankton communities and invasive species in the Berre Lagoon (Mediterranean Sea, France). *Marine Pollution Bulletin* 64: 1921–1932, <https://doi.org/10.1016/j.marpolbul.2012.06.020>
- Dorgham MM, Hamdy R (2015) The role of alien polychaetes along the Alexandria coast, Egypt. *International Journal of Environmental Research* 9: 141–150, <https://doi.org/10.22059/ijer.2015.883>
- Dyrynda PEJ, Fairall VR, Occhipinti-Ambrogi A, d'Hondt J-L (2000) The distribution, origins and taxonomy of *Tricellaria inopinata* d'Hondt and Occhipinti Ambrogi, 1985, an invasive bryozoan new to the Atlantic. *Journal of Natural History* 34: 1993–2006, <https://doi.org/10.1080/00222930050144828>
- Espinosa-Pérez MC, Hendrickx ME (2002) Distribution and ecological notes on *Dynoides* (Isopoda: Sphaeromatidae) in the Mexican Pacific. *Revista Brasileira de Biologia* 50: 643–647, http://www.scielo.sa.cr/scielo.php?script=sci_arttext&pid=S0034-77442002000200021&lng=en&nrm=iso
- Essl F, Bacher S, Genovesi P, Hulme PE, Jeschke JM, Katsanevakis S, Kowarik I, Kühn I, Pyšek P, Rabitsch W, Schindler S (2018) Which taxa are alien? Criteria, applications, and uncertainties. *BioScience* 68: 496–509, <https://doi.org/10.1093/biosci/biy057>
- Essl F, Dullinger S, Genovesi P, Hulme PE, Jeschke JM, Katsanevakis S, Kühn I, Lezner B, Pauchard A, Pyšek P, Rabitsch W, Richardson DM, Seebens H, van Kleunen M, van der Putten WH, Vilà M, Bacher S (2019) A conceptual framework for range-expanding species that track human-induced environmental change. *BioScience* 69: 908–919, <https://doi.org/10.1093/biosci/biz101>
- Eyuboglu O (2022) Measures and gap analysis on the impact of non-indigenous species on the Black Sea ecosystem. *Pakistan Journal of Zoology* 53: 1419–1429, <https://doi.org/10.17582/journal.pjz/20200726170735>
- Ferrario J, Rosso A, Marchini A, Occhipinti-Ambrogi A (2018) Mediterranean non-indigenous bryozoans: an update and knowledge gaps. *Biodiversity and Conservation* 27: 2783–2794, <https://doi.org/10.1007/s10531-018-1566-2>
- Fleminger A, Hendrix Kramer S (1988) Recent introductions of an Asian estuarine copepod, *Pseudodiaptomus marinus* (Copepoda: Calanoida), in southern California embayments. *Marine Biology* 98: 535–541, <https://doi.org/10.1007/BF00391545>
- Fortič A, Al-Sheik Rasheed R, Almajid Z, Badreddine A, Báez JC, Belmonte-Gallegos A, Bettoso N, Borne D, Camisa F, Caracciolo D, Çinar ME, Crocetta F, Četković I, Doğan A, Galiya M, García de los Ríos y los Huertos A, Grech D, Guallart J, Gündeğer G, Kahrčić A, Karachle PK, Kulijer D, Lombarte A, Marković O, Martínez Jiménez E, Sukran Okudan E, Orlando-Bonaca M, Sartoretto S, Spinelli A, Tuney Kizilkaya I, Virgili R (2023) New records of introduced species in the Mediterranean Sea. *Mediterranean Marine Science* 24: 182–202, <https://doi.org/10.12681/mms.34016>
- Fortič A, Mavrič B (2018) First record of the bryozoan *Tricellaria inopinata* (d'Hondt & Occhipinti Ambrogi, 1985) from the Slovenian Sea. *Annals for Istrian and Mediterranean Studies, Series Historia Naturalis* 28: 155–160, <https://doi.org/10.19233/ASHN.2018.19>
- Galanidi M, Aissi M, Ali M, Bakalem A, Bariche M, Bartolo AG, Bazairi H, Beqiraj S, Bilecenoglu M, Bitar G, Bugeja M, Carbonell A, Castriota L, Chalabi A, Çinar ME, Dragičević B, Dulčić J, El-Hawet AEA, Farrag MMS, Evans J, Galil B, Guerin L, Hyams-Kaphzan O, Kapedani R, Kamberi E, Livi S, Mačić V, Masse C, Mavrič B, Orlando-Bonaca M, Ouerghi A, Petovic S, Png-Gonzalez L, Schembri PJ, Shenkar N, Sghaier YR, Shakman E, Yahayaoui A, Yokeş MB, Zenetos A (2023) Validated Inventories of Non-Indigenous Species (NIS) for the

- Mediterranean Sea as Tools for Regional Policy and Patterns of NIS Spread. *Diversity* 15: 962, <https://doi.org/10.3390/d15090962>
- Galanidi M, Gratsia E, Zenetos A (2025) Backdating first records of non-indigenous species in the Mediterranean: are initial findings underestimated? *Mediterranean Marine Science* 26: 642–656, <https://doi.org/10.12681/mms.39662>
- Galil B (2008) *Sphaeroma walkeri* Stebbing, 1905 (Crustacea: Isopoda: Sphaeromatidae) established on the Mediterranean coast of Israel. *Aquatic Invasions* 3: 443–444, <https://doi.org/10.3391/ai.2008.3.4.13>
- Galil BS, Marchini A, Occhipinti-Ambrogi A, Minchin D, Narščiū A, Ojaveer H, Olenin S (2014) International arrivals: widespread bioinvasions in European seas. *Ethology Ecology & Evolution* 26: 152–171, <https://doi.org/10.1080/03949370.2014.897651>
- Gavira-O'Neill K, Guerra-García JM, Moreira J, Ros M (2016) Mobile epifauna of the invasive bryozoan *Tricellaria inopinata*: is there a potential invasional meltdown? *Marine Biodiversity* 48: 1169–1178, <https://doi.org/10.1007/s12526-016-0563-5>
- Gordon DR (1998) Effects of invasive, non-indigenous plant species on ecosystem processes: lessons from Florida. *Ecological Applications* 8: 975–989, [https://doi.org/10.1890/1051-0761\(1998\)008\[0975:EOINIP\]2.0.CO;2](https://doi.org/10.1890/1051-0761(1998)008[0975:EOINIP]2.0.CO;2)
- Grossi L, Bertasi F, Trabucco B (2017) New records of the alien polychaete worm *Chaetozona corona* (Polychaeta: Cirratulidae) in the Adriatic Sea. *Acta Adriatica* 58: 235–244, <https://doi.org/10.32582/aa.58.2.4>
- Harrison K, Holdich DM (1984) *Hemibranchiate sphaeromatids* (Crustacea: Isopoda) from Queensland, Australia, with a world-wide review of the genera discussed. *Zoological Journal of the Linnean Society* 81: 275–387, <https://doi.org/10.1111/j.1096-3642.1984.tb01175.x>
- Herdman WA (1906) Report on the Tunicata collected by Prof. Herdman at Ceylon in 1902. *Report to the Government of Ceylon on the Pearl Oyster Fisheries of the Gulf of Manaar* 5: 295–348
- Hidaka K, Itoh H, Hirai J, Tsuda A (2016) Occurrence of the *Paracalanus parvus* species complex in offshore waters south of Japan and their genetic and morphological identification to species. *Plankton and Benthos Research* 11: 131–143, <https://doi.org/10.3800/pbr.11.131>
- Hubendick B (1944) Beschreibung neuer Arten innerhalb der Gattung *Siphonaria*. *Arkiv för Zoologi* 35A(1): 1–7
- Jenkins B, Köhler F (2024) Hidden in plain sight: systematic review of Indo-West Pacific Siphonariidae uncovers extensive cryptic diversity (Mollusca, Gastropoda). *Megataxa* 13: 1–217, <https://doi.org/10.11646/megataxa.13.1.1>
- Johnson CH, Winston JE, Woollacott RM (2012) Western Atlantic introduction and persistence of the marine bryozoan *Tricellaria inopinata*. *Aquatic Invasions* 7: 295–303, <https://doi.org/10.3391/ai.2012.7.3.001>
- Karako S, Perl-Treves R, Katcoff D (2002) *Asterina burtoni* (Asteroidea; Echinodermata) in the Mediterranean and the Red Sea: does asexual reproduction facilitate colonization? *Marine Ecology Progress Series* 234: 139–145, <https://doi.org/10.3354/meps234139>
- Katsanevakis S, Wallentinus I, Zenetos A, Leppäkoski E, Çınar ME, Öztürk B, Grabowski M, Golani D, Cardoso AC (2014) Impacts of invasive alien marine species on ecosystem services and biodiversity: a pan-European review. *Aquatic Invasions* 9: 391–423, <https://doi.org/10.3391/ai.2014.9.4.01>
- Katsanevakis S, Poursanidis D, Hoffman R, Rizgalla J, Rothman SB-S, Levitt-Barmats Y, Hadjioannou L, Trkov D, Garmendia JM, Rizzo M, Bartolo AG, Bariche M, Tomas F, Kleitou P, Schembri PJ, Kletou D, Tiralongo F, Pergent C, Pergent G, Azzurro E, Bilecenoglu M, Lodola A, Ballesteros E, Gerovasileiou V, Verlaque M, Occhipinti-Ambrogi A, Kytinou E, Dailianis T, Ferrario J, Crocetta F, Jimenez C, Evans J, Ragkousis M, Lipej L, Borg JA, Dimitriadis C, Chatzigeorgiou G, Albano PG, Kalogirou S, Bazairi H, Espinosa F, Ben Souissi J, Tsiamis K, Badalamenti F, Langeneck J, Noel P, Deidun A, Marchini A, Skouradakis G, Royo L, Sini M, Bianchi CN, Sghaier Y-R, Ghanem R, Doumpas N, Zaouali J, Tsirintanis K, Papadakis O, Morri C, Çınar ME, Terrados J, Insacco G, Zava B, SoufiKechaou E, Piazzi L, Ounifi Ben Amor K, Andriotis E, Gambi MC, Ben Amor M M, Garrabou J, Linares C, Fortić A, Digenis M, Cebrian E, Fourt M, Zotou M, Castriota L, Di Martino V, Rosso A, Pipitone C, Falautano M, García M, Zakhama-Sraieb R, Khamassi F, Mannino AM, Ktari MH, Kosma I, Rifi M, K arachle P K, Yapıcı S, Bos AR, Balistreri P, Ramos Esplá AA, Tempesti J, Inglese O, Giovos I, Damalas D, Benhisoune S, Huseyinoglu MF, RjibaBahri W, Santamaria J, Orlando-Bonaca M, Izquierdo A, Stamouli C, Montefalcone M, Cerim H, Golo R, Tsioli S, Orfanidis S, Michailidis N, Gaglioti M, Taşkın E, Mancuso E, Žunec A, Cvitković I, Filiz H, Sanfilippo R, Siapatis A, Mavrič B, Karaa S, Türker A, Monniot F, Verdura J, El Ouamari N, Selfati M, Zenetos A (2020) Unpublished Mediterranean records of marine alien and cryptogenic species. *BioInvasions Records* 9: 165–182, <https://doi.org/10.3391/bir.2020.9.2.01>
- Katsanevakis S, Zaiko A, Olenin S, Costello MJ, Gallardo B, Tricarico E, Adriaens T, Jeschke JM, Sini M, Burke N, Ellinas K, Rutten S, Poursanidis D, Marchini A, Brys R, Raeymaekers JAM, Noé N, Hermoso V, Blaailid R, Lucy FE, Verbrugge LNH, Staehr PAU, Vandepitte L, de Groot D, Elliott M, Reuver M, Maclaren J, Li M, Oldoni D, Mazaris A, Trygonis V, Hablützel PI, Everts T, Pistevos JCA, Dekeyser S, Kimmig SE, Rickowski FS, Panov VE (2024) GuardIAS – Guarding European Waters from Invasive Alien Species. *Management of Biological Invasions* 15: 701–730, <https://doi.org/10.3391/mbi.2024.15.4.14>

- Kazmi QB, Yousuf F (2013) Checklist of Peracarida of Pakistan A—Marine and B— Terrestrial Isopoda marine Tanaidacea. *Fuast Journal of Biology* 3(2): 129–130
- Khalaji-Pirbalouty V, Gagnon JM (2021) A new species of *Dynoides* Barnard, 1914 (Crustacea, Isopoda, Sphaeromatidae) from Canada, with notes on geographic distribution of the north-eastern Pacific Ocean species. *Marine Biology Research* 17: 12–20, <https://doi.org/10.1080/17451000.2021.1892766>
- Khelifi-Touhami M, Ounissi M (2023) *Paracalanus* Boeck, 1864. In: ICES Identification Leaflets for Plankton. Vol. 199. ICES, Copenhagen, pp 1–22
- Kideys AE, Kovalev AV, Shulman G, Gordina A, Bingel F (2000) A review of zooplankton investigations of the Black Sea over the last decade. *Journal of Marine Systems* 24: 355–371, [https://doi.org/10.1016/S0924-7963\(99\)00095-0](https://doi.org/10.1016/S0924-7963(99)00095-0)
- Kondilatos G, Corsini Foka M, Pancucci Papadopoulou MA (2010) Occurrence of the first nonindigenous ascidian *Phallusia nigra* Savigny, 1816 (Tunicata: Ascidiacea) in Greek waters. *Aquatic Invasions* 5: 181–184, <https://doi.org/10.3391/ai.2010.5.2.08>
- Koukouras A, Sinis AI, Bobori D, Kazantzidis S, Kitsos M-S (2007) The echinoderm fauna of the Aegean Sea, and comparison with those of neighbouring seas. *Journal of Biological Research* 7: 67–92
- Kousteni V, Tsiamis K, Gervasini E, Zenetos A, Karachle PK, Cardoso AC (2022) Citizen scientists contributing to alien species detection: the case of fishes and molluscs in European marine waters. *Ecosphere* 13: e03875, <https://doi.org/10.1002/ecs2.3875>
- Kurt Şahin G, Çınar ME (2009) Eunicidae (Polychaeta) species in and around Iskenderun Bay (Levantine Sea, Eastern Mediterranean) with a new alien species for the Mediterranean Sea and a re-description of *Lysidice collaris*. *Turkish Journal of Zoology* 33: 331–347. <https://doi.org/10.3906/zoo-0806-19>
- Kurt-Sahin G, Çınar ME (2017) Distribution of Eunicidae (Annelida: Polychaeta) along the Levantine coast of Turkey, with special emphasis on alien species. *Marine Biodiversity* 47: 421–431, <https://doi.org/10.1007/s12526-016-0483-4>
- Langeneck J, Lezzi M, Del Pasqua M, Musco L, Gambi MC, Castelli A, Giangrande A (2020) Non-indigenous polychaetes along the coasts of Italy: a critical review. *Mediterranean Marine Science* 21: 238–275, <https://doi.org/10.12681/mms.21860>
- Langeneck J, Putignano M, Dimichele D, Giangrande A, Bilan M, Toso A, Musco L (2024) Non-indigenous polychaetes along the Salento Peninsula: new records and first molecular data. *Mediterranean Marine Science* 25: 184–203, <https://doi.org/10.12681/mms.35851>
- Latsoudis P (2020) A most northern record of alien tunicate *Phallusia nigra* Savigny, 1816, population from the Argolic Gulf, Greece. *Natural and Engineering Sciences* 5: 192–197, <https://doi.org/10.28978/nesciences.832997>
- Le Garrec V, Grall J, Chevalier C, Guyonnet B, Jourde J, Lavesque N, Bonifácio P, Blake JA (2017) *Chaetozona corona* (Polychaeta, Cirratulidae) in the Bay of Biscay: a new alien species for the Northeast Atlantic waters? *Journal of the Marine Biological Association of the United Kingdom* 97: 433–445, <https://doi.org/10.1017/S0025315416000540>
- Li L (2000) A new species of *Dynoides* (Crustacea: Isopoda: Sphaeromatidae) from the Cape d'Aguilar Marine Reserve, Hong Kong. *Australian Museum* 52: 137–149, <https://doi.org/10.3853/j.0067-1975.52.2000.1311>
- Liang D, Uye S (1997) Population dynamics and production of the planktonic copepods in a eutrophic inlet of the Inland Sea of Japan. *Marine Biology* 128: 415–421
- Link H, Nishi E, Tanaka K, Bastida-Zavala R, Kupriyanova EK, Yamakita T (2009) *Hydroides dianthus* (Polychaeta: Serpulidae), and alien species introduced into Tokyo Bay, Japan. *Marine Biodiversity Records* 2: 1–5, <https://doi.org/10.1017/S1755267209000931>
- Lodola A, Savini D, Occhipinti Ambrogi A (2012) First record of *Tricellaria inopinata* (Bryozoa: Candidae) in the harbours of La Spezia and Olbia, Western Mediterranean Sea (Italy). *Marine Biodiversity Records* 5: e41, <https://doi.org/10.1017/S1755267212000309>
- López de la Cuadra CM, García-Gómez JC (1994) Zoogeographical study of the Cheilostomatida from the Straits of Gibraltar. In: Hayward PJ, Ryland JS, Taylor PD (eds), *Biology and Palaeobiology of Bryozoans*, Olsen & Olsen, Fredensborg, Denmark, pp 107–112
- Lundberg B (1980) Selectivity of food algae by the herbivorous fish *Siganus rivulatus* in the marine vegetation at Mikhmoret (the Mediterranean coast of Israel). PhD Thesis, Hebrew University, 176 pp
- Magliozzi C, Cardoso AC, Son MO, Kvach Y, Japoshvili B, Stefanova K, Kurt G, Pantea E, Tabarcea C, Torodova V, Varshanidze M (2025) Data towards assessing non-indigenous species introductions in the Black Sea. *Marine Policy* 178: 106696
- Marin IN, Antokhina TI (2022) A new symbiotic scale worm (Polychaeta: Polynoidae) living in association with burrowing callinassid shrimps in the Black Sea. *Zoosystematica Rossica* 31: 272–285, <https://doi.org/10.31610/zsr/2022.31.2.272>
- Martin D, Britayev TA (1998) Symbiotic polychaetes: review of known species. *Oceanography and Marine Biology an Annual Review* 36: 217–340
- Martínez-Ortí A, Nappo A, Escutia V (2020) Nuevos hallazgos de los gasterópodos *Aplous assimilis* (Reeve, 1846) (F. Pisaniidae) y *Mitrella psilla* (Duclos, 1846) (F. Columbidae) en la costa mediterránea española. *Arxius de Miscel·lània Zoològica* 18: 51–57, <https://doi.org/10.32800/amz.2020.18.0051>
- Massé C, Viard F, Humbert S, Antajan E, Auby I, Bachelet G, Bernard G, Bouchet VMP, Burel T, Dauvin J-C, Delegrange A, Derrien-Courtet S, Droual G, Gouillieux B, Goulletquer P, Guérin L, Janson AL, Jourde J, Labruné C, Lavesque N, Leclerc JC, Le Duff M, Le Garrec V, Noël P,

- Nowaczyk A, Pergent-Martini C, Pezy JP, Raoux A, Raybaud V, Ruitton S, Sauriau PG, Spilmont N, Thibault D, Vincent D, Curd A (2023) An overview of marine non-indigenous species found in three contrasting biogeographic metropolitan French regions: insights on distribution, origins and pathways of introduction. *Diversity* 15: 161, <https://doi.org/10.3390/d15020161>
- Mastrototaro F, Brunetti R (2006) The non-indigenous ascidian *Distaplia bermudensis* in the Mediterranean: comparison with the native species *Distaplia magnilarva* and *Distaplia lucillae* sp. nov. *Journal of the Marine Biological Association of the United Kingdom* 86: 181–185, <https://doi.org/10.1017/S0025315406013014>
- Mastrototaro F, Montesanto F, Salonna M, Grieco F, Trainito E, Chimienti G, Gissi C (2019) Hitchhikers of the sea: concurrent morphological and molecular identification of *Symplesma brakenhielmi* (Tunicata: Ascidiacea) in the western Mediterranean Sea. *Mediterranean Marine Science* 20: 197–207, <https://doi.org/10.12681/mms.19390>
- Matishov G, Sohin I, Bulysheva N, Kovalenko M (2022) Expansion of oriental shrimp *Palaemon macrodactylus* Rathbun, 1902 (Crustacea, Decapoda, Palaemonidae) in the Azov-Don Basin. *Russian Journal of Biological Invasions* 15: 108–113, <https://doi.org/10.35885/1996-1499-15-3-108-113>
- Mayhoub H (1989) Envahissement de la côte Syrienne par une algue brune de mer Rouge. *Damascus University Journal* 5(18): 65–79
- Melki S, Ben Ahmed R, Gammoudi M, Azzouna A (2026) Rapid assessment investigation of ascidian species in five localities in the north of Tunisia: new record of three non-indigenous species *Polyclinum constellatum* Savigny, 1816, *Aplidium* cf. *accarense* (Millar, 1953), and *Polyandrocarpa zorritensis* (Van Name, 1931). *Marine Biodiversity* 56: 3, <https://doi.org/10.1007/s12526-025-01610-4>
- Micael J, Jardim N, Núñez C, Occhipinti-Ambrogi A, Costa AC (2016) Some Bryozoa species recently introduced into the Azores: reproductive strategies as a proxy for further spread. *Helgoland Marine Research* 70: 7, <https://hmr.biomedcentral.com/articles/10.1186/s10152-016-0458-7>
- Micu D, Nita V (2009) First record of the Asian prawn *Palaemon macrodactylus* Rathbun, 1902 (Caridea: Palaemonoidea: Palaemonidae) from the Black Sea. *Aquatic Invasions* 4: 597–604, <https://doi.org/10.3391/ai.2009.4.4.5>
- Mnasri-Afifi I, Zribi I, Abdelkader A, Charfi-Cheikhrouha F, Zakhama-Sraieb R (2024a) Rapid assessment survey of non-indigenous and cryptogenic species in Tunisian marinas. *BioInvasions Records* 13: 419–436, <https://doi.org/10.3391/bir.2024.13.2.10>
- Mnasri-Afifi I, Djebbi S, Zribi I, Naccache C, Charfi-Cheikhrouha F, Mezghanu Khemakhem M, Zakhama-Sraieb R (2024b) Unveiling the Hidden Identities of *Botrylloides niger* Herdman, 1886 in Tunisian Marinas Using DNA Barcoding. *Pakistan Journal of Zoology* 57: 2421–2428, <https://doi.org/10.17582/journal.pjz/20230505110532>
- Molnar JL, Gamboa RL, Revenga C, Spalding MD (2008) Assessing the global threat of invasive species to marine biodiversity. *Frontiers in the Ecology & the Environment* 6: 458–492, <https://doi.org/10.1890/070064>
- Montesanto F, Chimienti G, Gissi C, Mastrototaro F (2021) Spread of the non-indigenous ascidian *Aplidium accarense* (Millar, 1953) in the Eastern Mediterranean Sea: morphological and molecular tools for an accurate identification. *Mediterranean Marine Science* 22: 246–254, <https://doi.org/10.12681/mms.24887>
- Montesanto F, Chimienti G, Gissi C, Mastrototaro F (2022) *Polyclinum constellatum* (Tunicata, Ascidiacea), an emerging non-indigenous species of the Mediterranean Sea: Integrated taxonomy and the importance of reliable DNA barcode data. *Mediterranean Marine Science* 23: 121–135, <https://doi.org/10.12681/mms.28311>
- Munari C, Bocchi N, Parrella P, Granata T, Moruzzi L, Massara F, De Donati M, Mistri M (2017) The occurrence of two morphologically similar *Chaetozone* (Annelida: Polychaeta: Cirratulidae) species from the Italian seas: *Chaetozone corona* Berkeley & Berkeley, 1941 and *C. carpenteri* McIntosh, 1911. *The European Zoological Journal* 84: 541–553, <https://doi.org/10.1080/24750263.2017.1393843>
- Mureşan M, Teacă A, Begun T (2021) First record of the hydrozoan *Podocorynoides minima* (Trinci, 1903) in the Romanian Black Sea waters. *Biologia* 77: 1819–1828, <https://doi.org/10.1007/s11756-022-01051-5>
- Mytilineou Ch, Akel ESHK, Babali N, Balistreri M, Bariche M, Boyaci YÖ, Cilenti L, Constantinou C, Crocetta F, Çelik M, Dereli H, Dounas F, Durucan A, Garrido A, Gerovasileiou V, Kapiris K, Kebapcioglu T, Kleitou P, Krystalas A, Lipej L, Maina I, Marakis P, Mavrič B, Moussa R, Peña-Rivas L, Poursanidis D, Renada W, Rizkalla SI, Rosso A, Scirocco T, Sciuto F, Servello G, Tiralongo F, Yapici S, Zenetos A (2016) New Mediterranean Biodiversity Records (November 2016). *Mediterranean Marine Science* 17: 794–821, <https://doi.org/10.12681/mms.1976>
- Nappo A, Strizzi MNT, Mancini E, Marcelli M (2019) First record of *Mitrella psilla* (Duclos, 1846) (Gastropoda: Columbellidae) in Italy. *Bollettino Malacologico* 55: 62–64
- Nicolaidou A, Alongi G, Aydogan O, Catra M, Cavas L, Cevik A, Dosi A, Cirosta V, Giakoumi S, Gimenez Casaldueiro F, Filiz H, Izquierdo-Munoz A, Kalogirou S, Konstantinidis E, Kousteni V, Kout J, Legaki A, Megalofonou P, Ovalis P, Paolillo G, Paschos I, Perdikaris C, Poursanidis D, Ramos-Espla AA, Reizopoulou S, Sperone E, Taskin E, Triperi S, Vazquez-Luis M (2012) New Mediterranean Biodiversity Records (June 2012). *Mediterranean Marine Science* 13: 36–48, <https://doi.org/10.12681/mms.33>

- Nizamuddin M, Godeh M (1989) *Stypopodium tubruquense* (Phaeophyta, Dictyotales) a new species from the Mediterranean Sea. *Willdenowia* 18: 603–608
- Nunomura N (2010) A new species of the genus *Dynoides* (Isopoda: Sphaeromatidae) from Shiretoko, Hokkaido, northern Japan. *Contributions from the Toyama Science Museum* 384: 65–69
- Obst M, Exter K, Allcock AL, Arvanitidis C, Axberg A, Bustamante M, Cancio I, Carreira-Flores D, Chatzinikolaou E, Chatzigeorgiou G, Christmas N, Clark MS, Comtet T, Dailianis T, Davies N, Deneudt K, de Cerio OD, Fortić A, Gerovasileiou V, Hablützel PI, Keklikoglou K, Kotoulas G, Lasota R, Leite BR, Loisel S, Lévêque L, Levy L, Malachowicz M, Mavrić B, Meyer CA, Mortelmans J, Norkko J, Pade N, Power AM, Ramšak A, Reiss H, Solbakken J, Staehr PA, Sundberg P, Thyrring J, Troncoso JS, Viard F, Wenne R, Yperifanou EI, Zbawicka M, Pavludi C (2020) A Marine Biodiversity Observation Network for Genetic Monitoring of Hard-Bottom Communities (ARMS-MBON). *Frontiers in Marine Science* 7: 572680, <https://doi.org/10.3389/fmars.2020.572680>
- Occhipinti Ambrogio A (2000) Biotic invasions in a Mediterranean Lagoon. *Biological Invasions* 2: 165–176, <https://doi.org/10.1023/A:1010004926405>
- Occhipinti Ambrogio A, Marchini A, Cantone G, Castelli A, Chimenz C, Cormaci M, Frogia C, Furnari G, Gambi MC, Giaccone G, Giangrande A, Gravili C, Mastrototaro F, Mazziotti C, Orsi Relini L, Piraino S (2011) Alien species along the Italian coasts: an overview. *Biological Invasions* 13: 215–237, <https://doi.org/10.1007/s10530-010-9803-y>
- Omer-Cooper J (1927) Zoological results of the Cambridge Expedition to the Suez Canal, 1924, XII: report on the Crustacea Tanaidacea and Isopoda. *Transactions of the Zoological Society of London* 22: 201–209, <https://doi.org/10.1111/j.1096-3642.1927.tb00332.x>
- Orfanidis S, Alvito A, Azzurro E, Badreddine A, Ben Souissi J, Chamorro M, Crocetta F, Dalyan C, Fortić A, Galanti L, Geyran K, Ghanem R, Goruppi A, Grech D, Katsanevakis S, Madrenas E, Mastrototaro F, Montesanto F, Pavičić M, Pica D, Pola L, Pontes M, Ragkousis M, Rosso A, Sánchez-Tocino L, Tierno de Figueroa JM, Tiralongo F, Tirelli V, Tsioli S, Tunçer S, Vrdoljak D, Vuletin V, Zaouali J, Zenetos A (2021) New alien Mediterranean biodiversity records. *Mediterranean Marine Science* 22: 206–227, <https://doi.org/10.12681/mms.25294>
- Orsi JJ, Walter J, Chad T (1991) *Pseudodiaptomus forbesi* and *P. marinus* (Copepoda: Calanoida), the latest copepod immigrants to California's Sacramento-San Joaquin estuary. In: Proceedings 4th International Conference on Copepoda. *Bulletin of the Plankton Society of Japan* Special volume: 553–562
- Ounifi-Ben Amor K, Rifi M, Ghanem R, Draeif I, Zaouali J, Ben Souissi J (2015) Update of alien fauna and new records from Tunisian marine waters. *Mediterranean Marine Science* 17: 124, <https://doi.org/10.3906/zoo-1405-78>
- Peleg O, Guy-Haim T, Yeruham E, Silverman J, Rilov G (2020) Tropicalization may invert trophic state and carbon budget of shallow temperate rocky reefs. *Journal of Ecology* 108: 844–854, <https://doi.org/10.1111/1365-2745.13329>
- Péres JM (1958) Ascidies récoltées sur les côtes Méditerranéennes d'Israël. *Bulletin of the Research Council of Israel* 7B: 143–150
- Peters J, Albaina A, Blanco-Bercial L, Bucklin A, Di Capua I, Dos Santos A, Falkenhaus T, Fernandez de Puelles M, Fileman E, Garić R, Holst S, Hosia A, Laakmann S, Majaneva S, Maňko MK, Mazzocchi MG, O'Brien T, Peijnenburg KTCA, Peters A, Renz J, Rossel S, Weydmann-Zwolicka A, Winkler G, Yebra L, Cornils A (2025). Taxonomic uncertainty in North Atlantic and Mediterranean zooplankton limits species-level monitoring accuracy. *ICES Journal of Marine Science* 82: fsaf077, <https://doi.org/10.1093/icesjms/fsaf077>
- Ragkousis M, Zenetos A, Ben Souissi J, Hoffman R, Ghanem R, Taşkın E, Muresan M, Karpova E, Slynko E, Dağlı E, Fortić A, Surugiu V, Mačić V, Trkov D, Rjiba Bahri W, Tsiamis K, Ramos-Espla AA, Petrović S, Ferrario J, Marchini A, Sconfiati R, Ammar I, Alo A, Edelist D, Begun T, Teaca A, Tari G, Huseyinoglu MF, Karachle PK, Dogrammatzi A, Apostolopoulos GA, Crocetta F, Kytinou E, Digenis M, Skouradakis G, Tomas F, Bariche M, Kaminas A, Konida K, Deidun A, Marrone A, Fraschetti S, Mihneva V, Bianchi CN, Morri C, Gerovasileiou V, Lipej L, Sini M, Mangialajo L, Zotou M, Skolka M, Azzurro E, Vella A, Dailianis T, Grigoriou P, Jimenez C, Tsirintanis K, Oikonomidis G, Mancini E, Papadakis O, Di Martino V, Chatzigeorgiou G, Ben Amor MM, Vernadou E, Arda Y, Minasidis V, Azzola A, Hadjioannou L, Montefalcone M, Baldacchino Y, Stancanelli B, Bonifazi A, Occhipinti-Ambrogio A, Smeraldo S, Evans J, Kondylatos G, Falautano M, Castriota L, Lamprou A, Rizgalla J, Mavrić B, Papadimitrou E, Kersting D, Schembri PJ, Khamassi F, Nikolaou A, Ballesteros E, Dimitriadis C, Garcia M, Anastasiadis A, Kalogirou S, Nalmpanti M, Altamirano M, Grech D, Mavrouleas D, Vella N, Agius Darmanin S, Dragičević B, Poursanidis D, Tsatiris A, Corsini-Foka M, Orlando-Bonaca M, Insacco G, Tsalapatis A, Scannella D, Tiralongo F, Verdura J, Vitale S, Valsamidis MA, Bazairi H, Mannino AM, Virgili R, Collepardo Coccia F, El Zrelli R, Nikolidakis S, Jiani Rabaui L, Yapici S, Zaouali J, Zava B, Agrotis N, Bilecenoglu M, Çinar ME, Moraitis ML, Albano PG, Kaddouri N, Kosma I, Falsone F, Fossati V, Geraci M, Zamuda LL, Mancuso FP, Petrou A, Resaikos V, Aydın I, Batjakas IE, Bos AR, El Ouamari N, Giallongo G, Kampouris TE, Ounifi-Ben Amor K, Doğan A, Dulčić J, Şükran Okudan E, Rilov G, Rosso A, Royo L, Selfati M, Gaglioti M, Giakoumi S, Kousteni V, Micu D, Nicoara M, Orfanidis S, Papatheodoulou M, Tempesti J, Triantaphyllou M, Tsourou T, Yalgin F, Baltag E, Cerim H, Filiz H, Gerogiadis CG, Papadamakis P, Rammou DL, Samargiu MD, Sciuto F, Sinopoli M, Türkler A, Chiarore A, Tamburello L, Karray S, Hassen B, Katsanevakis S (2023) Unpublished Mediterranean and Black Sea records of marine alien, cryptogenic, and neonative species. *BioInvasions Records* 12: 339–369, <https://doi.org/10.3391/bir.2023.12.2.01>

- Ramos-Esplá AA, Izquierdo-Muñoz A, Çınar ME (2013) *Microcosmus exasperatus* (Ascidacea, Pyuridae), current distribution in the Mediterranean Sea. *Marine Biodiversity Records* 6: e89, <https://doi.org/10.1017/S1755267213000663>
- Ramos-Esplá AA, Bitar G, Ramzi Sghaier Y, Çınar ME, Deidun A, Ferrario J, Ulman A (2020) *Symplesma* (Ascidacea: Styelidae), a non-indigenous genus spreading within the Mediterranean Sea: taxonomy, routes and vectors. *Aquatic Invasions* 15: 44–62, <https://doi.org/10.3391/ai.2020.15.1.04>
- Raykov V, Lepage M, Perez-Dominguez R (2010) First record of oriental shrimp, *Palaemon macrondactylus* Rathbun, 1902 in Varna Lake, Bulgaria. *Aquatic Invasions* 5: S91–S95, <https://doi.org/10.3391/ai.2010.5.S1.019>
- Reverter-Gil O, Souto J (2025) New and non-indigenous species of Bryozoa from Iberian waters. *European Journal of Taxonomy* 885: 33–64, <https://doi.org/10.5852/ejt.2023.885.2187>
- Reyes-Martínez JM, González-Gordillo JI (2019) New record of the non-indigenous copepod *Pseudodiaptomus marinus* Sato, 1913 (Calanoida, Pseudodiaptomidae) from the Guadalquivir estuary (Gulf of Cadiz, SW Spain). *Crustaceana* 92: 675–683, <https://doi.org/10.1163/15685403-00003903>
- Rosso A (1994) Segnalazione di *Electra tenella* (HINCKS) (Bryozoa) lungo il versante sud-orientale della Sicilia. *Bollettino dell'Accademia Gioenia di Scienze Naturali, Catania* 27: 241–251
- Rosso A (2009) The first cateniceid (Bryozoa, Asciophora) from Mediterranean shallow-waters: a hidden resident or a new immigrant? *Journal of Natural History* 43: 2209–2226, <https://doi.org/10.1080/00222930903089977>
- Rosso A, Di Martino E (2023) Capturing the moment: a snapshot of Mediterranean bryozoan diversity in the early 2023. *Mediterranean Marine Science* 24: 426–445, <https://doi.org/10.12681/mms.34329>
- Rosso A, Di Martino E, Pica D, Galanti L, Cerrano C, Novosel M (2018) Non-indigenous bryozoan species from natural and artificial substrata of Mediterranean submarine caves. *Marine Biodiversity* 48: 1345–1355, <https://doi.org/10.1007/s12526-016-0602-2>
- Rosso A, Sanfilippo R, Sciuto F, Serio D, Catra M, Alongi G, Viola A, Leonardi R (2019) Preliminary information on bryozoans associated with selected *Cystoseira* communities from Sicily (Mediterranean). *Memoirs of the Australasian Association of Palaeontologists* 52: 115–129
- Rousou M, Langeneck J, Apserou C, Arvanitidis C, Charalambous S, Chrysanthou K, Constantinides G, Dimitriou PD, García Gómez SC, Hadjieftychiou SI, Katsiaras N, Kleitou P, Kletou D, Küpper FC, Louizidou P, Martins R, Moraitis ML, Papageorgiou N, Papatheodoulou M, Petrou A, Xevgenos D, Vasiliades L, Voultsiadou E, Chinatioglou CC, Castelli A (2023) Polychaetes (Annelida) of Cyprus (Eastern Mediterranean Sea): an updated and annotated checklist including new distribution records. *Diversity* 15: 941, <https://doi.org/10.3390/d15080941>
- San Martín G, Lucas Y, Hutchings P (2024) The genus *Syllis* Savigny in Lamarck, 1818 (Annelida: Syllidae: Syllinae) from Australia (Fourth part). *Zootaxa* 5453: 1–32, <https://doi.org/10.11646/zootaxa.5453.1.1>
- Sartoni G, De Biasi AM (1999) A survey of the marine algae of Milos Island, Greece. *Cryptogamie Algologie* 20: 271–283, [https://doi.org/10.1016/S0181-1568\(99\)80019-1](https://doi.org/10.1016/S0181-1568(99)80019-1)
- Scuderi D, Villari A (2022) Further two alien species for the Sicilian waters: *Aplus assimilis* (Reeve, 1846) and *Mitrella psilla* (Duclos, 1846) (Gastropoda Neogastropoda). *Biodiversity Journal* 13: 355–359, <https://doi.org/10.31396/Biodiv.Jour.2022.13.2.355.359>
- Schroeder K, Chiggiato J, Bryden HL, Borghini M, Ben Ismail S (2016) Abrupt climate shift in the Western Mediterranean Sea. *Scientific Reports* 6: 23009, <https://doi.org/10.1038/srep23009>
- Seebens H, Schwartz N, Schupp PJ, Blasius B (2016) Predicting the spread of marine species introduced by global shipping. *Proceedings of the National Academy of Sciences* 113: 5646–5651, <https://doi.org/10.1073/pnas.1524427113>
- Shenkar N, Loya Y (2009) Non-indigenous ascidians (Chordata: Tunicata) along the Mediterranean coast of Israel. *Marine Biodiversity Records* 2: e166, <https://doi.org/10.1017/S1755267209990753>
- Simberloff D, Von Holle B (1999) Positive interactions of nonindigenous species: invasional meltdown? *Biological Invasions* 1: 21–32
- Streftaris N, Zenetos A, Papathanassiou E (2005) Globalisation in marine ecosystems: the story of non-indigenous marine species across European Seas. *Oceanography and Marine Biology* 43: 419–453, <https://doi.org/10.1201/9781420037449>
- Subías-Barata A, Sánchez-Vidal A, Di Martino E, Figuerola B (2022) Marine biofouling organisms on beached, buoyant and benthic plastic debris in the Catalan Sea. *Marine Pollution Bulletin* 175: 113405, <https://doi.org/10.1016/j.marpolbul.2022.113405>
- Sun R, Yang D (2000) Study on *Hydroides* (Polychaeta: Serpulidae) from waters of China. I. *Studia Marina Sinica* 42: 116–135
- Sun Y, Wong E, Keppel E, Williamson JE, Kupriyanova EK (2017) A global invader or a complex of regionally distributed species? Clarifying the status of an invasive calcareous tubeworm *Hydroides dianthus* (Verrill, 1873) (Polychaeta: Serpulidae) using DNA barcoding. *Marine Biology* 164: 28, <https://doi.org/10.1007/s00227-016-3058-9>
- Teacă A, Begun T, Muresan M (2021) Annelid invaders in the Black Sea region: The distribution of *Streblospio gynobranchiata* and first occurrence of *Laonome xeprovala*. *Global Ecology and Conservation* 32: e01920, <https://doi.org/10.1016/j.gecco.2021.e01920>
- Teacă A, Begun T, Menabit S, Muresan M (2022) The first record of *Marenzelleria neglecta* and the spread of *Laonome xeprovala* in the Danube Delta-Black Sea ecosystem. *Diversity* 14: 423, <https://doi.org/10.3390/d14060423>

- Temiz B, Öztürk E, Blanchoud S, Karahan A (2023) Phylogeographic and morphological analysis of *Botrylloides niger* Herdman, 1886 from the northeastern Mediterranean Sea. *Diversity* 15: 367, <https://doi.org/10.3390/d15030367>
- Terbiyik T, Cevik C, Toklu-Alicli B, Sarihan E (2007) First record of *Ferosagitta galerita* (Dallot, 1971) [Chaetognatha] in the Mediterranean Sea. *Journal of Plankton Research* 29: 721–726, <https://doi.org/10.1093/plankt/fbm053>
- Thessalou-Legaki E, Aydogan Ö, Bekas P, Bilge G, BoYaci YÖ, Brunelli E, Circosta V, Crocetta F, Durucan F, Erdem M, Ergolavou A, Filiz H, Fois F, Gouva E, Kapiris K, Katsanevakis S, Kljajić Z, Konstantinidis E, Konstantinou G, Koutsogiannopoulos D, Lamon S, Mačić V, Mazzete R, Meloni D, Mureddu A, Paschos I, Perdikaris C, Piras F, Poursanidis D, Ramos-Esplá AA, Rosso A, Sordino P, Sperone E, Sterioti A, Taskin E, Toscano F, Tripepi S, Tsiakkios L, Zenetos A (2012) New Mediterranean Biodiversity Records (December 2012). *Mediterranean Marine Science* 13: 312–327, <https://doi.org/10.12681/mms.313>
- Thomsen MS, Wernberg T, South PM, Schiel DR (2016) To include or not to include (the invader in community analyses)? That is the question. *Biological Invasions* 18: 1515–1521, <https://doi.org/10.1007/s10530-016-1102-9>
- Timofeev V, Simakova U, Spiridonov V (2019) The first finding of the oriental shrimp *Palaemon macrodactylus* Rathbun, 1902 (Crustacea: Decapoda, Palaemonidae) in the territorial waters of Russia in the Azov–Black Sea basin. *Russian Journal of Biological Invasions* 10: 192–198, <https://doi.org/10.1134/S2075111719020139>
- Tokioka T (1961) Ascidians collected during the Melanesia expedition of the Osaka Museum of Natural History. 1. Ascidians presented by Dr. R.I.A. Catala of the aquarium of Noumea. *Publications of the Seto Marine Biology Laboratory* 9: 103–138, <https://doi.org/10.5134/174659>
- Tortonese E (1966) Echinoderms from the coast of the Lebanon. *Miscellaneous papers on the Natural Sciences, in the American University of Beirut* 5: 2–5
- Toso A, Putignano M, Fumarola LM, Bariche M, Giangrande A, Musco L, Piraino S, Langeneck J (2024) A revised inventory of Annelida in the Lebanese coastal waters with ten new aliens for the Mediterranean Sea. *Mediterranean Marine Science* 25: 715–731, <https://doi.org/10.12681/mms.37998>
- Tsirintanis K, Azzurro E, Crocetta F, Dimiza M, Frogia C, Gerovasileiou V, Langeneck J, Mancinelli G, Rosso A, Stern N, Triantaphyllou M, Tsiamis K, Turon X, Verlaque M, Zenetos A, Katsanevakis S (2022) Bioinvasion impacts on biodiversity, ecosystem services, and human health in the Mediterranean Sea. *Aquatic Invasions* 17: 308–352, <https://doi.org/10.3391/ai.2022.17.3.01>
- Tsiamis K, Panayotidis P, Zenetos A (2008) Alien marine macrophytes in Greece: a review. *Botanica Marina* 51: 237–246, <https://doi.org/10.1515/BOT.2008.040>
- Tsiamis K, Zenetos A, Deriu I, Gervasini E, Cardoso AC (2018) The native distribution range of the European marine non-indigenous species. *Aquatic Invasions* 13: 187–198, <https://doi.org/10.3391/ai.2018.13.2.01>
- Tsiamis K, Palialexis A, Stefanova K, Gladan ŽN, Skejić S, Despalatović M, Cvitković I, Dragičević B, Dulčić J, Vidjak O, Bojanić N (2019) Non-indigenous species refined national baseline inventories: A synthesis in the context of the European Union's Marine Strategy Framework Directive. *Marine Pollution Bulletin* 145: 429–435, <https://doi.org/10.1016/j.marpolbul.2019.06.012>
- Tsiamis K, Palialexis A, Connor D, Antoniadis S, Bartilotti C, Bartolo AG, Berggreen-Ulrik C, Boschetti S, Buschbaum C, Canning-Clode J, Carbonell A, Castriota L, Corbeau C, Costa A, Cvitković I, Despalatović M, Dragičević B, Dulčić J, Fortić A, Francé J, Gittenberger A, Gizzi F, Gollasch S, Gruszka P, Hegarty M, Hema T, Jensen K, Josephides M, Kabuta S, Kerckhof F, Kovtun-Kante A, Krakau M, Kraśniewski W, Lackschewitz D, Lehtiniemi M, Lieberum C, Linnamägi M, Lipej L, Livi S, Lundgreen K, Magliozzi C, Massé C, Mavrić B, Michailidis N, Moncheva S, Mozetič P, Naddafi R, Ninčević Gladan Ž, Ojaveer H, Olenin S, Orlando-Bonaca M, Ouerghi A, Parente M, Pavlova P, Peterlin M, Pitacco V, Png-Gonzalez L, Rousou M, Sala-Pérez M, Serrano A, Skorupski J, Smolders S, Srebaliene G, Stæhr PA, Stefanova K, Strake S, Tabarcea C, Todorova V, Trkov D, Tuaty-Guerra M, Vidjak O, Žuljević A, Cardoso AC (2021) Marine Strategy Framework Directive Descriptor 2, Non-Indigenous Species: delivering solid recommendations for setting threshold values for non-indigenous species pressure on European seas. Publications Office of the European Union, 22 pp, <https://doi.org/10.2760/035071>
- Turon X, Nishikawa T, Rius M (2007) Spread of *Microcosmus squamiger* (Ascidacea: Pyuridae) in the Mediterranean Sea and adjacent waters. *Journal of Experimental Marine Biology and Ecology* 342: 185–188, <https://doi.org/10.1016/j.jembe.2006.10.040>
- Tzomos T, Chartosia N, Christodoulou M, Kitsos M-S (2011) New records and range expansion of lessepsian migrants in the Levantine and Aegean Seas. *Marine Biodiversity Records* 3: e111, <https://doi.org/10.1017/S1755267209991114>
- Ulman A, Ferrario J, Occhipinti-Ambrogi A, Arvanitidis C, Bandi A, Bertolino M, Bogi C, Chatzigeorgiou G, Çiçek BA, Deidun A, Ramos-Esplá A, Koçak C, Lorenti M, Martinez-Laiz G, Merlo G, Princisgh E, Scribano G, Marchini A (2017) A massive update of non-indigenous species records in Mediterranean marinas. *PeerJ* 5: e3954, <https://doi.org/10.7717/peerj.3954>
- Uttieri M, Anadoli O, Banchi E, Battuello M, Beşiktepe Ş, Carotenuto Y, Marques SC, de Olazabal A, Di Capua I, Engell-Sørensen K, Goruppi A, Guy-Haim T, Hure M, Kourkoutmani P, Lučić D, Mazzocchi MG, Michaloudi E, Morov AR, Kurt TT, Tirelli V, Vannini J, Velassquez X, Vidjak O, Wootton M (2023) The distribution of *Pseudodiaptomus*

- marinus* in European and neighbouring waters—a rolling review. *Journal of Marine Science and Engineering* 11: 1238, <https://doi.org/10.3390/jmse11061238>
- Uye SI, Iwai Y, Kasahara S (1982) Reproductive biology of *Pseudodiaptomus marinus* (Copepoda: Calanoida) in the Inland Sea of Japan. *Bulletin of the Plankton Society of Japan* 29(1): 25–35
- Varrella S, Livi S, Corinaldesi C, Castriota L, Maggio T, Vivona P, Pindo M, Fava S, Danovaro R, Dell’Anno A (2025) A comprehensive assessment of non-indigenous species requires the combination of multi-marker eDNA metabarcoding with classical taxonomic identification. *Environment International* 199: 109489, <https://doi.org/10.1016/j.envint.2025.109489>
- Verlaque M, Boudouresque CF (1991) *Stypopodium schimperi* (Buchinger ex Kützing) Verlaque et Boudouresque comb. nov. (Dictyotales, Fucophyceae), algue de mer Rouge récemment apparue en Méditerranée. *Cryptogamie, Algologie* 12: 195–211, <https://doi.org/10.5962/p.309317>
- Verlaque M, Ruitton S, Mineur F, Boudouresque CF (2015) CIESM Atlas of exotic species in the Mediterranean. Vol. 4 *Macrophytes*. CIESM Publishers, Monaco, 364 pp
- Virgili R, Tanduo V, Katsanevakis S, Terlizzi F, Villani G, Fontana A, Crocetta F (2022) The Miseno Lake (central-western Mediterranean Sea): an overlooked reservoir of non-indigenous and cryptogenic ascidians in a marine reserve. *Frontiers in Marine Science* 9: 866906, <https://doi.org/10.3389/fmars.2022.866906>
- Virgili R, Fasciglione P, Tiralongo F, Pannone R, Palomba M, Santoro M, De Luca P, Crocetta F (2025) Risk assessment at early invasion stages: interdisciplinary approaches maximise the evidence on human-induced and climate-driven fish dispersal. *Hydrobiologia* 852: 2241–2261, <https://doi.org/10.1007/s10750-024-05537-y>
- Walter TC (1989) Review of the New World Species of *Pseudodiaptomus* (Copepoda: Calanoida), with a key to the species. *Bulletin of Marine Science* 45(3): 590–628
- Zenetos A, Çinar ME, Crocetta F, Golani D, Rosso A, Servello G, Shenkar N, Turon X, Verlaque M (2017) Uncertainties and validation of alien species catalogues: The Mediterranean as an example. *Estuarine, Coastal and Shelf Science* 191: 171–187, <https://doi.org/10.1016/j.ecss.2017.03.031>
- Zenetos A, Galanidi M (2020) Mediterranean non indigenous species at the start of the 2020s: recent changes. *Marine Biodiversity Records* 13: 10, <https://doi.org/10.1186/s41200-020-00191-4>
- Zenetos A, Gofas S, Russo G, Templado J (2004) CIESM Atlas of exotic species in the Mediterranean. Vol. 3 *Molluscs*. CIESM Publishers, Monaco, 380 pp
- Zenetos A, Gofas S, Verlaque M, Çinar ME, García Raso JE, Bianchi CN, Morri C, Azzurro E, Bilecenoglu M, Frogia C, Siokou I, Violanti D, Sfriso A, San Martín G, Giangrande A, Katağan T, Ballesteros E, Ramos-Esplá A, Mastrototaro F, Ocaña O, Zingone A, Gambi MC, Streftaris N (2010) Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union’s Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Mediterranean Marine Science* 11: 381–493, <https://doi.org/10.12681/mms.87>
- Zenetos A, Gofas S, Morri C, Rosso A, Violanti D, Garcia Raso J, Cinar M, Almogi-Labin A, Ates A, Azzurro E, Ballesteros E, Bianchi C, Bilecenoglu M, Gambi M, Giangrande A, Gravili C, Hyams-Kaphzan O, Karachle PK, Katsanevakis S, Lipej L, Mastrototaro F, Mineur F, Pancucci-Papadopoulou M, Ramos Espla A, Salas C, San Martin G, Sfriso A, Streftaris N, Verlaque M (2012) Alien species in the Mediterranean Sea by 2012. A contribution to the application of the European Union’s Marine Strategy Framework Directive (MSFD). Part 2. Introduction trends and pathways. *Mediterranean Marine Science* 13: 328–352, <https://doi.org/10.12681/mms.327>
- Zenetos A, Gratsia E, Cardoso A-C, Tsiamis K (2019) Time lags in reporting of biological invasions: the case of Mediterranean Sea. *Mediterranean Marine Science* 20: 469–475, <https://doi.org/10.12681/mms.20716>
- Zenetos A, Karachle PK, Corsini-Foka M, Gerovasileiou V, Simbora N, Xentidis NJ, Tsiamis K (2020) Is the trend in new introductions of marine non-indigenous species a reliable criterion for assessing good environmental status? The case study of Greece. *Mediterranean Marine Science* 21: 775–787, <https://doi.org/10.12681/mms.25136>
- Zenetos A, Tsiamis K, Galanidi M, Carvalho N, Bartilotti C, Canning-Clode J, Castriota L, Chainho P, Comas González R, Costa AC, Dragičević B, Dulčić J, Faasse M, Florin AB, Gittenberger A, Jakobsen H, Jelmert A, Kerckhof F, Lehtiniemi M, Livi S, Lundgreen K, Macic V, Massé C, Mavrič B, Naddafi R, Orlando-Bonaca M, Petovic MS, Png-Gonzalez L, Carbonel Quetglas A, Ribeiro RS, Cidade T, Smolders S, Stæhr PAU, Viard V, Outinen O (2022a) Status and trends in the rate of introduction of marine non-indigenous species in European seas. *Diversity* 14: 1077, <https://doi.org/10.3390/d14121077>
- Zenetos A, Albano PG, Lopez Garcia E, Stern N, Tsiamis K, Galanidi M (2022b) Established non-indigenous species increased by 40% in 11 years in the Mediterranean Sea. *Mediterranean Marine Science* 23: 196–212, <https://doi.org/10.12681/mms.29106>
- Zibrowius H (1973) Remarques sur trois espèces de Serpulidae acclimatées en Méditerranée: *Hydroides dianthus* (Verrill, 1873), *Hydroides dirampha* Mörch, 1863, et *Hydroides elegans* (Haswell, 1883). *Rapport de la Commission Internationale de la Mer Méditerranée* 21: 683–686
- Zibrowius H, Bitar G (2003) Invertébrés marins exotiques sur la côte du Liban. *Lebanese Science Journal* 4: 67–74
- Žuljević A, Lučić P, Cvitković I, Despalatović M (2023) Marine brown algae *Stypopodium schimperi* is going to be the most invasive alien species in the Adriatic Sea. In: Jelaska S. 5th Croatian Symposium on Invasive species with International Participation book of abstracts, Croatia Ecological Society, Zagreb, pp 31–31

Websites, online databases and software

- Alvite N (2023) Characterization of the zooplanktonic community and detection of introduced species in ports using DNA metabarcoding. [Master's Thesis, University of A Coruña]. <https://www.udc.es/es/tfe/traballo/?codigo=30917> (accessed 1 June 2025)
- Bitar G, Dupuy de la Grandrive R, Foulquié M (2003) Second mission relating to the development of marine protected areas in Syrian coasts, pp 1–18. http://medmpa.rac-spa.org/pdf/Rapports/Syrie/Mission%20Report_October%202003 (accessed 01 June 2024)
- Boyko CB, Campos-Filho IS, Hadfield KA, Hughes T, Merrin KL, Ota Y, Poore GCB (eds) (2025) World Marine, Freshwater and Terrestrial Isopod Crustaceans database. *Dynoides* Barnard, 1914. <https://www.marinespecies.org/Isopoda/aphia.php?p=taxdetails&id=204371> (accessed 1 June 2025)
- Muñoz IA, Ramos-Esplá AA (2019) Updating the list of recent non-indigenous ascidians (Chordata: Tunicata) and its spreading in the Mediterranean Sea. Ten years later (2009–2019). *Frontiers in Marine Science*. Conference Abstract: XX Iberian Symposium on Marine Biology Studies (SIEBM XX). <https://doi.org/10.3389/conf.fmars.2019.08.00188> (accessed 5 May 2020)
- WoRMS Editorial Board (2025) World Register of Marine Species. <https://www.marinespecies.org>. (accessed 30 May 2025)

Supplementary material

The following supplementary material is available for this article:

Table S1. Dataset of unpublished Mediterranean and Black Sea records of marine alien, cryptogenic, and neonative species (III)
http://www.reabic.net/journals/bir/2026/Supplements/BIR_2026_Pistevos_etal_SupplementaryTable.xlsx

Figure S1. *Leodice antennata* from Lebanon

Figure S2. *Syllis ergeni* from Algeria

Figure S3. *Syllis schulzi* from Cyprus

Figure S4. *Prionospio depauperata* from Piombino, Tyrrhenian Sea

Figure S5. *Chaetozone corona* from the Gulf of Trieste

Figure S6. *Hydroides dianthus* from Romania

Figure S7. *Dynoides amblysinus* from Port of Catania, Italy

Figure S8. *Sphaeroma walkeri* from Syria

Figure S9. *Mesanthura cf. romulea* from Algeria

Figure S10. *Palaemon macrodactylus* zoea from Romania

Figure S11. *Pseudodiaptomus marinus* found in Algeciras Harbour (Spain)

Figure S12. *Paracalanus tropicus* found in Algeciras Harbour (Spain)

Figure S13. *Siphonaria crenata* from Syria

Figure S14. *Mitrella psilla* from Algeria

Figure S15. (A) *Catenicella paradoxa* from Sicily Strait (B) Articulated biserial internodes of *Tricellaria inopinata* from Sicily (C) *Arbopercula angulata* from Liguria

Figure S16. (A, B) Regenerated specimens of *Aquilonastra burtoni* recorded from Syria

Figure S17. *Ferosagitta* sp. from the Black Sea

Figure S18. (A, B) *Symplegma rubra* off the coast of Tunisia

Figure S19. *Symplegma brakenhielmi* from Alicante port (Western Mediterranean). Chromomorphs: red (A) and yellow (B)

Figure S20. *Phallusia nigra* from Italy

Figure S21. *Microcosmus exasperatus* from Italy

Figure S22. *Distaplia bermudensis* (A) Bizerte, Tunisia; (B) Brindisi, Italy; (C) El Kantaoui, Tunisia

Figure S23. *Botrylloides niger* colony from Greece

Figure S24. *Stypopodium schimperi* in Lampedusa Island

http://www.reabic.net/journals/bir/2026/Supplements/BIR_2026_Pistevos_etal_SupplementaryFigures.pdf